

Description
of the
Demographic and Health Surveys
Individual Recode
Data File

MEASURE *DHS* +

Version 1.0

(with differences from DHS III)

March 5, 2008

Foreword

DHS surveys collect primary data using several types of questionnaires. A household questionnaire is used to collect information on characteristics of the household's dwelling unit, and data related to the height and weight for women and children in the household. It is also used to identify members of the household who are eligible for an individual interview. Eligible respondents are then interviewed using an individual questionnaire.

In a majority of DHS surveys eligible individuals include women of reproductive age (15-49) and men age 15-59, or in some cases 15-54. In some countries only women are interviewed. Individual questionnaires include information on fertility, family planning and maternal and child health. Data are available from DHS for each of these surveys by request through the mail or from our web site at www.measuredhs.com. Data from DHS surveys are produced in both raw and recode formats. A raw data file includes the data as they were collected, without any structural changes. These files are generally not distributed, but they are also available on request. A recode data file is in a standardized format, with the same structure across countries participating in each DHS phase. This standardization is meant to facilitate comparisons across surveys. This document describes the standard recode defined for the fourth round of DHS surveys (MEASURE *DHS+*). Recode structures are defined for households, women and men.

DHS also collects data using other types of surveys and questionnaires. These include surveys of education, health service providers, communities, household health expenditures, young adults, and others. These data are also available, but there are no recode definitions for them.

Data Archive,
Demographic and Health Surveys - MEASURE *DHS+*

Table of Contents

General Description	1
Introduction.....	1
Rationale for Recoding	1
Data File Structure	2
Coding Standards	3
Respondent Identification	4
Record Identification	4
Survey Identification.....	4
Century Month Code.....	5
Imputed Dates	5
Model Questionnaires	6
Sections and Occurrences	6
Section and Variable Descriptions.....	6
Section and Variable Description	6
Section and Variable Description - Household.....	6
Section and Variable Description - Men.....	6

General Description

Introduction

This document contains two parts. The first part is a general discussion of the recode file, including the rationale for recoding; description of the physical structure in which the recode file is available; coding standards used in the data file; location of identification information; use of century month codes for dates and imputation of partial dates; DHS model questionnaires; sections and occurrences. The second part provides a description of each variable in the data file, giving additional information that is not available in the dictionary.

Rationale for Recoding

The individual data are transformed into a standardized recode dataset for several reasons:

- First, dates for several key events are imputed as much analysis of the data is based on these events and their dates are often incomplete or missing. The imputed dates are included in the data file to allow analysts to produce results consistent with those published by DHS and to save analysts the time and trouble of creating their own imputation schemes.
- Second, variables as collected in the original questionnaire are in a form convenient for collection but not always for analysis. Often the same question is asked in several places in the questionnaire, but to different respondents. In the recode file these variables are combined and created in a form that is easy to use for analysis.
- Third, summary variables are often necessary in analysis and many of these, including the summary variables that are used in the DHS reports, are included in the recode file.
- Fourth, certain indices, particularly the anthropometric indices from the height and weight data, are calculated from the data and included in the recode file.
- Finally, and in many ways most importantly, the data in the recode file are in a standardized format allowing easy comparison of data between countries.

The DHS approach to creating standardized individual recode data files for each country is part of the DHS policy to make the data accessible, providing the analyst with the data in the most convenient form for analysis. This approach, while providing easy access to the data, is not without its pitfalls. **DHS strongly suggests that analysts become familiar with the questionnaires used in the surveys they are analyzing.** The questionnaires used in one country, while containing essentially the same information, may be different in many ways from those used in another country. In creating the standardized individual recode data files these differences require special consideration and total standardization is obviously not possible. The recode data file is structured in two parts, standard sections and country-specific sections. The standard sections contain the same variables in the same positions for all countries. The country-specific sections contain all variables specific to the country and so are not standardized across countries.

Data File Structure

The recode data file is available in three different structures; the structure to use depends on the hardware and software requirements of the analyst:

- Flat** Each record of the data file represents one case (respondent), with all variables being placed one after the other on the same record. The repeating sections of the recode file are placed one after the other on the record, with the maximum number of occurrences of each section being represented in the data file. Each variable in a repeating section is placed immediately after the preceding variable of the same occurrence, such that all variables for occurrence 1 precede all variables for occurrence 2 of a section. For example, in the birth history BIDX, BORD, B0, B1 etc. for the first occurrence appear followed by the second occurrence of BIDX, BORD, B0, B1 etc. The length of the records in the data file are fixed, exceeding 4000 characters in total. The total size of the data file is on average approximately 40M bytes, depending on the sample size, with the largest files being over 380 M bytes in size. The flat file is designed for users using statistical packages that only support data structures containing a fixed number of records per case. This format is similar to the format of the World Fertility Survey standard recode files. An SPSS/PC+, SAS or STATA data file description is distributed with this file format.
- Rectangular** Each case (respondent) in the data file contains a fixed number of records, with each record representing a section of the data file. For repeating sections there is a record for each occurrence of the section, with the maximum number of occurrences of each repeating section being included in the data file. The number of records in a data file will vary from country to country as the number of country-specific sections of the data file varies, but for the standard sections of the data file there are 70 records, excluding the calendar. For data files distributed on magnetic tape the record length of each record will be fixed at the length of the longest record in the data file, but for PC users the record length will vary, with each record terminating with a CR/LF, as for standard DOS text files. The total size of the data file is about 55 M bytes, with the largest files being over 550 M bytes in size. On magnetic tape the file size is considerably larger. The rectangular file is designed for microcomputer users using software that requires a fixed number of records per case. An SPSS/PC+ , SAS or STATA data file description is distributed with this file format.
- Hierarchical** The hierarchical data structure is identical to the rectangular data structure, with the exception that records exist only for the occurrences of the sections that are necessary. As an example of the difference, if a woman has 6 children there will be 6 records in the birth history section in the hierarchical structure, but 20 records (the maximum number of occurrences for this section) in the rectangular data structure, with the last 14 occurrences filled with blanks. The record length will be the same as for the rectangular file. The total size of the file is approximately 25 M bytes, depending on the sample size, with the largest files being over 265 M bytes in size. The hierarchical data structure is designed for use with ISSA, the Integrated System for Survey Analysis, available from DHS and CPro. An ISSA dictionary is distributed with this file format.

Coding Standards

Special codes are used throughout the data file for certain responses. The general coding scheme is presented below. The codes given apply to 4 digit, 3 digit, 2 digit and 1 digit variables, respectively. If there are other special responses to questions, these are coded in decreasing order from these special codes, i.e., 9996, 996, 96, 6; 9995, 995, 95, 5; etc.

BLANK	Variable is <u>not applicable</u> for this respondent either because the question was not asked in a particular country or because the question was not asked of this respondent due to the flow or skip pattern of the questionnaire.
9999, 999, 99, 9	This question should have been answered by the respondent, but the questionnaire contained no information for this variable (<u>missing data</u>).
9998, 998, 98, 8	The respondent replied " <u>Don't know</u> " to this question.
9997, 997, 97, 7	The answer to this question was <u>inconsistent</u> with other responses in the questionnaire and it was thought that this response was probably in error. The response was changed to this code to avoid further problems due to inconsistency of information. This usually takes place during the secondary editing stage of data processing.

In addition a code of 0 is generally used as a negative response in the data file. For example, "No education" is coded 0 for V106, "No problem" is coded 0 for V467A, and a simple response of "No" is coded 0 in all standard sections of the data file. In the country-specific sections of the data file, variables are generally coded in the same way as they were on the questionnaire and a "No" answer usually has code 0.

In certain questions a two-digit coding scheme is used in which the first digit, representing the major coding category, is standard, but the second digit is country-specific. This applies to questions such as those relating to water source, toilet facilities, and source of contraception. For example, for source of contraception the major categories are:

- 1 Public Sector
- 2 Private Medical Sector
- 3 Other Private Sector
- 4 Other

The coding scheme for V326 (last source of contraception for current users of modern methods) might use codes such as:

- 11 Government hospital
- 12 Government health center
- ...
- 21 Private hospital or clinic
- 22 Private doctor
- ...
- 31 Shop
- ...

In the above coding scheme, the first digit is the standard major category; the second digit is country-specific.

Respondent Identification

Each record of the data file starts with the identification for each case in the data file, and has the variable name CASEID (see description of CASEID). It occupies the first 15 character positions of each record, irrespective of the type of data file structure.

Record Identification

For rectangular and hierarchical data files, each record has an identifying code in character positions 16-17 of the record. This record identification identifies the section of the data file that is contained on the record (e.g., 21 for the birth history). Repeating sections will have the same record identification for each occurrence of the section. In the hierarchical data files a variable following the record identification in each section specifies which occurrence of the section the record represents; in the rectangular data files the variable exists for all occurrences that are non-blank.

Survey Identification

For each survey there is a two-character alphabetic country identification code plus a one-digit data structure code in variable V000. The variable V000 occupies positions 16-18 of the record for flat files, and positions 18-20 of the first record of the rectangular or hierarchical data files. The one-digit data structure code is always 3 4 for MEASURE *DHS III*+ surveys, except for those MEASURE *DHS* + surveys that used DHS III Model questionnaires. The country codes are as follows:

DHS IV:

Armenia	AM	Gabon	GA	Malawi	MW	Rwanda	RW
Bangladesh	BD	Ghana	GH	Mali	ML	Tanzania	TZ
Benin	BJ	Guatemala	GU	Mauritania	MR	Turkey	TR
Cambodia	KH	Guinea	GN	Namibia	NM	Turkmenistan	TM
Colombia	CO	Haiti	HT	Nepal	NP	Uganda	UG
Dominican R.	DR	Jordan	JO	Nicaragua	NC	Zambia	ZM
Egypt	EG	India	IA	Nigeria	NG	Zimbabwe	ZW
Ethiopia	ET	Kazakhstan	KK	Peru	PE		

Century Month Code

All dates in the data file are expressed in terms of months and years and also as century month codes. A century month code (CMC) is the number of the month since the start of the century. For example, January 1900 is CMC 1, January 1901 is CMC 13, January 1980 is CMC 961, September 1994 is CMC 1137. The CMC for a date is calculated from the month and year as follows:

$$\text{CMC} = (\text{YY} * 12) + \text{MM} \quad \text{for month MM in year 19YY.}$$

To calculate the month and year from the CMC use the following formulae:

$$\text{YY} = \text{int}((\text{CMC} - 1) / 12)$$

$$\text{MM} = \text{CMC} - (\text{YY} * 12)$$

For Dates in 2000 and after the CMC is calculated as follows:

$$\text{CMC} = ((\text{YYYY}-1900) * 12) + \text{MM} \quad \text{for month MM in year YYYY.}$$

To calculate the month and year from the CMC use the following formulae:

$$\text{YYYY} = \text{int}((\text{CMC} - 1) / 12) + 1900$$

$$\text{MM} = \text{CMC} - ((\text{YYYY}-1900) * 12)$$

Imputed Dates

For key events in the respondent's life, dates have been imputed when the full date of the event was not provided by the respondent or in some cases if dates are inconsistent (e.g. less than 7 months between births). These events are the date of birth of the respondent, the date of first union or marriage, the date of birth of each child of the respondent, the date of conception of the current pregnancy (based on the duration of pregnancy), the date of start of use of current method, and. For each of these dates only the imputed data are available in the recode data file, but a date flag has been included in the file to show what format the information was in prior to imputation, and what basis was used for the imputation. The codes for this date flag are as follows:

- 1 Both month and year of the event were specified and so no imputation was necessary.
- 2 The year of the event was not given, but the month of the event and the age of the respondent or child or, in the case of the date of first union, the respondent's age at first union were specified. In most cases this information uniquely identifies the exact date of the event. In a few cases the year of the event was imputed from a choice of two possible years.
- 3 The year of the event, but not the month, and the age of the respondent or child or, in the case of the date of first union, the respondent's age at first union were specified and only the month of the event was imputed.
- 4 The year of birth, but not the month, and the age of the respondent or child were specified. However, in surveys where it is believed the year of birth is calculated from the age, the year of birth is ignored when the year of birth plus the age add up to the year of interview.
- 5 The year of the event was given but the month of the event was not specified, and neither was the age. The month of the event was imputed.
- 6 Neither the month nor the year of the event were specified, but age was given and the year and month of the event were imputed from the age.
- 7 Only the month of the event was given, without the year or age. The year of the event was imputed from other information. (For current pregnancy, duration of pregnancy was given.)

- 8 No information was given concerning the date of the event. But month and year of the event were imputed from other information. (For current pregnancy, duration of pregnancy was not given.)

For the date of conception of the current pregnancy only codes 7 and 8 are used. The date of interview is required to be fully specified in all cases and so no imputation is necessary for this variable and no format flag exists for the date of interview.

A full description of the imputation process is given in the DHS Data Processing Manual.

Model Questionnaires

Two core questionnaires were used during the DHS surveys, Model "A" questionnaire for High Contraceptive Prevalence Countries and Model "B" questionnaire for Low Contraceptive Prevalence Countries. The two questionnaires contain basically the same information, although the Model "A" questionnaire contains a detailed calendar of events in the five years preceding the interview, whereas the Model "B" questionnaire contains a simpler series of questions.

In the variable description section that follows, the column labeled "Model" indicates in which questionnaire the question is asked. An "A" indicates that the variable refers to a question asked only in countries that used a Model "A" questionnaire, and a "B" indicates that the variable relates to a question asked only in countries that used the Model "B" questionnaire. If the column is blank, then the question is asked in both Model "A" and Model "B" questionnaires. If the column contains an "X", then the question is not included in either of the Model questionnaires, but was used in a sufficient number of surveys to justify its inclusion as a standard variable. If the column contains "MM", then the questions come from the maternal mortality module. If the column contains "FG", then the questions come from the female genital cutting module.

Sections and Occurrences

The data file is broken down into a number of logical sections. These sections translate directly into records for the rectangular and hierarchical data structures. The logical sections are designed to map the sections of the model questionnaires, although some sections of the model questionnaire are split into more than one section in the recode data file. Some of these sections are repeating or multiple occurrence sections while others are single occurrence sections. Single sections contain simple, single-answer variables.

Multiple sections are used to represent sets of questions that are repeated for a number of events. The birth history is an example of a multiple section, where questions relating to children are asked for each child, and each child has an entry in the birth history. Each entry in the multiple section is known as an occurrence of the section. In rectangular and hierarchical data files each occurrence of the section occupies a separate record. Multiple sections are used for sets of questions where the number of occurrences may vary.

In contrast, sets of questions for which there are a fixed number of occurrences are held in a group. A group is similar to a multiple section, but is stored on a single record for rectangular and hierarchical files. In addition single variables may also be included in a section containing a group. In the recode file the contraceptive table (REC31) is stored as a group containing 20 entries, one for each contraceptive method. For the flat files there is no difference between groups and multiple sections.

Section and Variable Descriptions

The section description following gives an outline of the sections of the recode file and the types of information they contain. The description is based on the rectangular and hierarchical files. The section description gives the name of the section, the section code used to identify the section in the data file, the length of the record for that section, the section class (S for single and M for multiple), the minimum and maximum number of occurrences of the section in each case, and the section label.

The section description is followed by variable descriptions. . The variable descriptions provide additional background information relating to each variable.

Section and Variable Description

<u>Section</u>	<u>Code</u>	<u>Length</u>	<u>Class</u>	<u>Occurrences</u>		<u>Section label</u>
				<u>Min</u>	<u>Max</u>	
REC01	01	129	S	1	1	without group Respondent's basic data
REC11	11	106	S	0	1	without group Respondent's basic data
REC21	21	54	M	0	20	Reproduction (Birth history)
REC22	22	98	S	0	1	Reproduction
REC31	31	99	S	0	1	Contraception Table
REC32	32	127	S	0	1	Contraception knowledge & use
REC41	41	196	M	0	6	Maternity
REC42	42	187	S	0	1	Health and Breastfeeding
REC43	43	201	M	0	6	Health
REC44	44	91	M	0	6	Height and Weight
REC51	51	59	S	0	1	Marriage/Exposure
REC61	61	70	S	0	1	Fertility preferences
REC71	71	54	S	0	1	Partner's Characteristics
REC75	75	224	S	0	1	AIDS and Condom Use
REC81	81	37	S	0	1	Characteristics of Interview
REC82	82	98	M	0	9	Calendar (optional)
REC83	83	52	M	0	20	Maternal mortality (optional)
REC84	84	37	S	0	1	Maternal mortality (optional)
REC91	91	?	S	0	1	Country specific - Single variables
REC92	92	?	M	0	20	Country specific - births
REC94	94	?	M	0	6	Country specific - Maternity
REC95	95	?	M	0	6	Country specific - Health
REC96	96	?	M	0	6	Country specific - Height/Weight
REC97	97	?	M	0	?	Country-specific
REC98	98	?	M	0	?	Country-specific
REC99	99	?	M	0	?	Country-specific

? implies that the entry is country-specific

Section 01 (REC01)

Respondent's Basic Data

<u>Var</u>	<u>Model</u>	<u>Description</u>
CASEID		Case identification, used to uniquely identify each respondent. In most surveys this is constructed by concatenating the cluster or sample point number, the household number and the respondent's line number, but in some surveys this may be the questionnaire number taken from the front page of the questionnaire.
V000		Alphabetic country code to identify the survey from which the data were collected. The code is based on an international standard code. This variable is 3 characters in length, with the third character indicating the format of the recode file used for this survey. For all surveys in MEASURE <i>DHS</i> III + this code will be 4. For example: 5 BJ4 is Benin, KH4 is Cambodia, CO4 is Colombia.
V001		Cluster number is the number identifying the sample point as used during the fieldwork. This variable may be a composite of several variables in the questionnaire. If so, the original variables are included in REC91 as country-specific variables.
V002		Household number is the number identifying the household in which the respondent was interviewed, within the sample point. In some cases, this variable may be the combination of dwelling number and household number within dwelling. In these cases, the original variables are included as country-specific variables.
V003		Respondent's line number in the household schedule.
V004		Ultimate area unit is a number assigned to each sample point to identify the ultimate area units used in the collection of data. This variable is usually the same as the cluster number, but may be a sequentially numbered variable for samples with a more complicated structure.
V005		Sample weight is an 8 digit variable with 6 implied decimal places. To use the sample weight divide it by 1000000 before applying the weighting factor. All sample weights are normalized such that the weighted number of cases is identical to the unweighted number of cases when using the full dataset with no selection. This variable should be used to weight all tabulations produced using the data file. For self-weighting samples this variable is equal to 1000000.
V006		Month of interview.
V007		Year of interview. The number of digits used during this phase is changed from 2 digits to 4 digits
V008		Century month code of date of interview (see note on century month codes).
V009		Month of birth of respondent (see note on imputed dates).
V010		Year of birth of respondent (see note on imputed dates). The number of digits used during this phase is changed from 2 digits to 4 digits
V011		Century month code of date of birth of the respondent (see note on century month codes).

<u>Var</u>	<u>Model</u>	<u>Description</u>
V012		Current age in completed years is calculated from the century month code of the date of birth of the respondent (V011) and the century month code of the date of interview (V008). In a few cases the age in the data file will be different from that reported by the respondent when the respondent's birthday was in the month of interview, but she had not yet had her birthday. If the respondent correctly reported her age at her last birthday (and not her age at her next birthday) then the calculated age was rounded up from the reported age, to avoid inconsistencies between the age and the century month code for the birth.
V013		Current age in 5-year groups is produced by grouping V012.
V014		Completeness of information for the date of birth of the respondent (see note on imputed dates). Codes for DHSII MEASURE DHS+ are different from the codes used in DHS I.
V015		Result of individual interview. Code 1 represents a completed interview. For all other cases, only REC01 will exist in the data file. For flat and rectangular format data files, cases with a result code different than 1 are dropped from the file.
V016		Day of the month in which the interview took place.
V017		Century month code for the first month of the calendar. This is constant for all cases and is the century month code of January of the first year of the calendar.
V018	A	Row of calendar representing the month of interview. The calendar is numbered from 1 to 80, with month 80 being January of the first year of the calendar. This variable is coded 0 for incomplete interviews or for questionnaires using the Model "B" questionnaire.
V019	A	Records the length of the calendar to use for this case. V019 is equal to 80-V018+1. This variable is coded 0 for incomplete interviews or for questionnaires using the Model "B" questionnaire.
V019A		Number of calendar columns. In the model B questionnaire only 1 column is used and 4 columns in the model A questionnaire.
V020		The ever-married sample indicator is a constant for all cases in the data file. For all woman samples it is code 0, and for ever married samples it is code 1.
V021		Primary sampling unit is a number assigned to sample points to identify the primary sampling units for use in the calculation of sampling errors. This variable is usually the same as the cluster number and/or the ultimate area unit, but may differ if the sample design required a multistage selection process.
V022		Sample strata defines the pairings or groupings of primary sampling units used in the calculation of sampling errors when using the Taylor series expansion method (for example, with the package Clusters).
V023		Sample domain defines the basic geographic units within which the sample was designed. For example, if the sample was designed to be self-weighting within region, this variable would define those regions; if the sample was designed to be self-weighting within major urban areas, other urban areas and rural areas, this variable would define the major urban,

<u>Var</u>	<u>Model</u>	<u>Description</u>
		other urban and rural areas. If the sample is self-weighted at the national level, this variable is code 0.
V024		<i>De facto</i> region of residence. This is a copy of V101, added to this section to allow for analysis of completion rates by region.
V025		<i>De facto</i> type of place of residence. This is a copy of V102, added to this section to allow for analysis of completion rates by urban/rural residence.
V026		<i>De facto</i> place of residence is the type of place in which the respondent was interviewed. This is a copy of V134, added to this section to allow for analysis of completion rates by type of place of residence (Capital city, small town, town and countryside).
V027		Number of visits for the interview. This is a copy of V804.
V028		Interviewer identification code. Codes are country-specific. This variable occupies 3 digits. This is a copy of V805.
V029		Data entry keyer code. Codes are country-specific. This is a copy of V806.
V030		Field supervisor's code. Codes are country-specific. This variable uses 3 digits in MEASURE <i>DHS+</i> .
V031		Field editor's code. Codes are country-specific. This variable uses 3 digits in MEASURE <i>DHS+</i> .
V032		Office editor's code. Codes are country-specific.
V033		Ultimate area unit selection probability is the probability of selection of the ultimate area unit, ignoring the household selection. This variable can be used in conjunction with data for the sample point, such as service availability data.
V034	X	Line number of husband as recorded in the household schedule. This variable can be used, in conjunction with the cluster or sample point number and the household number to match the women's data with the husbands' data, to allow for the analysis of couples.
V040		Cluster altitude in meters. Used to adjust the anemia level.
V042	X	Household selection for hemoglobin.
V043	X	Selection for women's status module. This variable is set to not applicable in case there is no women's status module.
V044	X	Selection for domestic violence module. This variable is set to not applicable in case there is no domestic violence module.

Section 11 (REC11)

Respondent's Basic Data

<u>Var</u>	<u>Model</u>	<u>Description</u>
V101		<i>De facto</i> region of residence. Region in which the respondent was interviewed. Codes are country-specific. This variable is now two digits. For <i>de jure</i> region of residence, see V139.
V102		<i>De facto</i> type of place of residence. Type of place of residence where the respondent was interviewed as either urban or rural. Note that this is not the respondent's own categorization, but was created based on whether the cluster or sample point number is defined as urban or rural. See also V134. For <i>de jure</i> type of place of residence, see V140.
V103		Childhood place of residence is classified into city, town and countryside as reported by the respondent. In some countries, additional codes are used for capital/major cities (code 0) and for abroad (code 4).
V104		Number of years the respondent has lived in the village, town, or city where she was interviewed. Visitors to the community are coded 96.
V105		Type of place of previous residence is coded as for V103. In some countries, additional codes are used for capital/major cities (code 0) and for abroad (code 4). BASE: All respondents except those answering "Always" or "Visitor" to V104 (V104 \diamond 95 & V104 \diamond 96).
V106		Highest education level attended. This is a standardized variable providing level of education in the following categories: No education, Primary, Secondary, Higher. In some countries the educational system does not fit naturally within this scheme and a different categorization was used for the Final Report. In this case, this variable is constructed as accurately as possible from the country's own scheme and the variable used for the Final Report is included as a country-specific variable.
V107		Highest year of education gives the years of education completed at the level given in V106. BASE: All respondents except those answering "No education" or with missing data for V106 (V106 \diamond 0 & V106 \diamond 9).
V108		Literacy of the respondent. In many countries, respondents with secondary or higher levels of education are coded 1, "Reads easily." The exact criteria for this assumption is country-specific.
V109		Whether the respondent usually reads a newspaper or magazine at least once a week.
V110		Whether the respondent usually watches television every week.
V111	A	Whether the respondent usually listens to a radio every day.
V112	B	Whether the respondent usually listens to a radio every week.
V113		Major source of drinking water for members of the household. Individual codes are country-specific, but the major categories are standard.
V114	X	Major source of water for household use other than for drinking. Individual codes are country-specific, but the major categories are standard.

<u>Var</u>	<u>Model</u>	<u>Description</u>																		
V115		Time taken to get to the water source for drinking water. BASE: All respondents except those with drinking water either piped to, or available from a well in, the residence, yard or plot, or who use rainwater or bottled water (V113 < 11 & V113 < 21 & V113 < 41 & V113 < 61). The actual selection criteria is country-specific.																		
V116		Type of toilet facility in the household. Individual codes are country-specific, but the major categories are standard.																		
		Whether the household has:																		
V119		Electricity.																		
V120		A radio.																		
V121		A television.																		
V122		A refrigerator.																		
		Whether a member of the household has:																		
V123		A bicycle.																		
V124		A motorcycle.																		
V125		A car.																		
V127		Main material of the floor. Individual codes are country-specific, but the major categories are standard.																		
V128	X	Main material of the walls. Individual codes are country-specific, but the major categories are standard.																		
V129	X	Main material of the roof. Individual codes are country-specific, but the major categories are standard.																		
V130		Religion. Both the question and the codes are country-specific.																		
V131		Ethnicity. Both the question and the codes are country-specific.																		
V133		Education in single years. This variable is constructed from the educational level (V106) and the grade at that level (V107) as follows: <table border="0" style="margin-left: 40px;"> <tr> <td>V106</td> <td>=></td> <td>V133</td> </tr> <tr> <td>0</td> <td>=></td> <td>0</td> </tr> <tr> <td>1</td> <td>=></td> <td>V107</td> </tr> <tr> <td>2</td> <td>=></td> <td>V107+x</td> </tr> <tr> <td>3</td> <td>=></td> <td>V107+y</td> </tr> <tr> <td>9</td> <td>=></td> <td>99</td> </tr> </table> <p>x = years to complete primary education y = years to complete primary and secondary education where both x and y are country-specific.</p>	V106	=>	V133	0	=>	0	1	=>	V107	2	=>	V107+x	3	=>	V107+y	9	=>	99
V106	=>	V133																		
0	=>	0																		
1	=>	V107																		
2	=>	V107+x																		
3	=>	V107+y																		
9	=>	99																		
V134		<i>De facto</i> place of residence is the type of place in which the respondent was interviewed. Urban areas are classified into large cities (capital cities and cities with over 1 million population), small cities (population over 50,000), and towns (other urban areas), and all rural areas are assumed to be countryside. Note that this classification differs from that used in DHS I.																		
V135		Whether the respondent is a usual resident of the household or is just visiting the household. Responses of "Visitor" to V104 are visitors to the city, town or village where the interview took place, but V135 shows respondents who were visitors to the household.																		

<u>Var</u>	<u>Model</u>	<u>Description</u>
V136		Total number of household members is the number of usual residents plus the number of visitors who slept in the house the previous night that were listed in the household schedule.
V137		Number of children resident in the household and aged 5 and under. Visiting children are not included.
V138		Number of eligible women in the household. Eligible women are usually defined to be women aged 15-49 who slept in the household the previous night, irrespective of whether they usually reside in the household or are visiting the household. In some countries an ever-married sample is used for the individual interview, and so the eligibility criteria is further restricted to ever-married women.
V139		<i>De jure</i> region of usual residence. For <i>de facto</i> region of residence, see V101.
V140		<i>De jure</i> type of place of usual residence. For <i>de facto</i> type of place of residence, see V102.
V141		<i>De jure</i> place of residence. In most countries, no differentiation is made between large cities and small cities in this variable.
V142	X	Whether the same source of water is used for drinking water as for household water.
V144	A	Whether the respondent lived in one or more than one community since January 1987.
V148		Whether the respondent is still in school. In DHS III, these data are now taken directly from a question in the women's questionnaire. Women who are older than 24 years of age are coded 0, assuming that they are no longer in school.
V149		Educational achievement recodes the education of the respondent into the following categories: None, incomplete primary, complete primary, incomplete secondary, complete secondary, higher education. See related variables V106, V107, V133.
V150		Relationship to the head of the household. These data are taken from the household schedule.
V151		Sex of the head of the household.
V152		Age of the head of the household.
V153		Whether the household has a telephone.
V154		Main reason that the respondent stopped attending school. BASE: Women who are under the age of 25 and who are not attending school (V012 < 25 & V148 < 1).

Inflation factors for ever-married samples

Variables AWFAC TT to AWFAC TE are standard inflation factors to be applied to the denominators when using ever-married samples to produce estimates for all women. To produce these estimates for all women it is necessary to apply the inflation factors to account for the proportion of women who were never married. Each factor is stored in 5-digit variables, with two implied decimal places. A value of 00128 means an inflation factor of 1.28 should be applied to the individual case to allow for never-married women. This means that for every 100 ever-married women found in the household schedule of a particular age and with the same background characteristic, there are 128 women in total, i.e. 100 ever-married women plus 28 never-married women. These inflation factors are used in the calculation of fertility rates, median ages at first union and first birth, mean number of children ever born, and other all-woman-based estimates. Note that these inflation factors do not need to be used when the denominator for an estimate only includes women who have ever been married. Four standard factors are produced for ever-married samples. Additional factors may appear as country-specific inflation factors if they were calculated to produce tabulations for the final report of a particular country. Country-specific variables are located in REC91.

AWFACTT	All-woman factor for the total population.
AWFACTU	All-woman factor for the urban/rural breakdowns.
AWFACTR	All-woman factor for the regional breakdowns.
AWFACTE	All-woman factor for the educational breakdowns.
V155	Literacy. Whether a respondent who attended primary schooling can read a whole or part of a sentence showed. A respondent who attended secondary education or higher are coded 2 as well as respondent who could read a whole sentence.
V156	Ever participated in a literacy program or any other program that involves learning to read and write (not including primary school)
V157	Frequency of reading newspaper or magazine
V158	Frequency of listening to radio
V159	Frequency of watching television
V160	Toilet facilities shared with other households
V161	Type of cooking fuel
V162	Place for hand washing
<u>Items present in the household as observed by the interviewer</u>	
V163	Presence of water/tap in household
V164	Presence of soap/ash/other cleansing agent in household
V165	Presence of basin in household
V166	Results of salt iodine test

Section 21 (REC21)

Reproduction

The birth history contains up to 20 entries for births, and is ordered in reverse order such that the last birth is given first in the birth history and the first birth is given last. For respondents with more than 20 births, the birth history contains the last 19 births plus the first birth. However, all variables relating to intervals between births are calculated based on the actual births, and not just the births given in the birth history. The variable V224 contains the count of entries in the birth history, and is thus the index to the last entry in the birth history which contains the information relating to the first birth.

<u>Var</u>	<u>Model</u>	<u>Description</u>
BIDX		Birth history index numbers the entries in the birth history from 1 to n, where the nth birth is the first birth.
BORD		Birth order number gives the order in which the children were born and so is the reverse order from BIDX.
B0		Twin code gives an order number for each child of a multiple birth. Code 0 indicates a single birth, code 1-upwards give the number of the child. Twins are ordered in the birth history with the higher twin codes appearing before the lower twin codes. See the example of the birth history structure below.
B1		Month of birth of child (see note on imputed dates).
B2		Year of birth of child (see note on imputed dates). This variable now occupies 4 digits.
B3		Century month code for the date of birth of the child (see note on century month codes).
B4		Sex of child.
B5		Whether child was alive or dead at the time of interview.
B6		Age at death of the child as reported in the questionnaire. The first digit of the age at death gives the units in which it was reported: 1 - Days, 2 - Months, 3 - Years, 9 - Special responses. The last two digits give the age at death in those units. Age at death is usually reported in days if it was less than one month, in months if it was less than two years and otherwise in years. If the last two digits contain a value greater than 90 then this is a special response. For example, 298 means the age at death was a number of months, and the exact number was unknown, but lies between 1 and 23 months. BASE: Dead children (B5 = 0).
B7		Age at death of the child in completed months gives a calculated age at death from the reported information. If it was reported in days these are truncated to completed months, if reported in months these are used directly, but if reported in years then truncated years are used, i.e., 3 years becomes 36 months. For ages at death that were not specified, an age at death is imputed using a hot deck approach by taking the same age at death as the last child encountered of the same birth order in the data file. This variable is no longer truncated at 90 months and now occupies three digits. BASE: Dead children (B5 = 0).
B8		Current age of the child in single years for all living children. BASE: Living children (B5 = 1).

<u>Var</u>	<u>Model</u>	<u>Description</u>
B9		The person the child usually lives with. The Respondent is coded 0, father coded 1, other relatives coded 2, other people coded 3, and children aged 15 and over who were not asked who they live with are coded 4. Note that this coding is different from DHS I. BASE: Living children (B5 = 1).
B10		Completeness of information for the date of birth of the child (see note on imputed dates). Codes are different from the codes used in DHS I.
B11		Preceding birth interval is calculated as the difference in months between the current birth and the previous birth, counting twins as one birth. BASE: All births except the first birth and its twins.
B12		Succeeding birth interval is calculated as the difference in months between the current birth and the following birth, counting twins as one birth. BASE: All births except the last birth and its twins.
B13		Flag for age at death is coded as follows: 0 No flag 1 Age at death plus the date of birth would place the death after the interview 2 Age at death is less than the reported duration of breastfeeding 3 Age at death is less than the age the child was first given supplemental foods 4 Age at death is less than age the child was first breastfed 5 Age at death plus the date of birth would place the death before the last vaccination 6 Reported age at death is outside the range expected for the units given 7 Age at death was imputed, however the units were given 8 Age at death was imputed, no units were given BASE: Dead children (B5 = 0).
B14		Whether the interval between the birth and the previously reported birth was four or more years. This calculation is performed by the interviewer, purely in terms of years, ignoring the month of birth of each of the children. For example, a birth in April 1988 would be categorized as being four or more years after a birth in October 1984, even though there is actually less than four years between the births. BASE: Second and higher births (BORD > 1).
B15		Whether there were any other live births in the interval between the birth and the previously reported birth. Note: Variables B14 and B15 are questions used in the interview to try and ascertain if the respondent had omitted any live births while reporting her birth history. The birth history may have been modified during the interviewing process, and the responses may not now correspond to the interval between the birth and the preceding birth recorded in this section. The data have been included to help indicate cases where the respondent may have omitted live births while originally reporting the birth history. The omitted births should be included in this final version of the birth history.
B16		Child's line number in household.

Var Model Description

Example Birth History:

BIDX	BORD	B0	B1	B2	B3	B4	B5	B6	B7	B8	B9	B10	B11	B12	B13	B15	B16
1	6	2	2	2000	1202	1	1			1	0	1	28			0	6
2	5	1	2	2000	1202	2	0	301	15			1	28		6	0	
3	4	0	10	1997	1174	2	1			3	0	4	43	28		0	5
4	3	0	3	1994	1131	1	1			7	0	4	14	43		0	7
5	2	0	1	1993	1117	2	0	302	24			5	17	14	0	0	
6	1	0	8	1991	1100	1	0	212	12			5		17	0		

In this example there are six children, including a pair of twins. There are three boys and three girls. Two of the girls have died, one after one year and the other after two years and one boy who died after 12 months, the other living children live with their mother. Exact dates of birth were available only for the last birth and its twin. For the other births either the year only (code 5 in B10) , or year and age (code4 in B10) were available. The birth intervals are calculated between births excluding children of multiple births. For example the preceding interval for the first entry is the difference between the CMC for the first entry and the third entry because the second entry is a twin of the first entry.

Section 22 (REC22)**Reproduction**

<u>Var</u>	<u>Model</u>	<u>Description</u>
V201		Total number of children ever born. If there are fewer than twenty births then this is the same as V224 (Number of entries in the birth history), but if there are more than twenty births then this gives the full number, while V224 will be 20.
V202		Total number of sons living at home.
V203		Total number of daughters living at home.
V204		Total number of sons living away from home.
V205		Total number of daughters living away from home.
V206		Total number of sons who have died.
V207		Total number of daughters who have died. V201 is the sum of variables V202 to V207.
V208		Total number of births in the last five years is defined as all births in the months 0 to 59 prior to the month of interview, where month 0 is the month of interview.
V209		Total number of births in the past year is defined as all births in the months 0 to 12 (not 0 to 11) prior to the month of interview.
V210		Total number of births in the month of interview.
V211		Century month code of the date of first birth is the same as B3 (V224). BASE: All respondents with one or more births (V201 > 0).
V212		Age of the respondent at first birth is calculated from the CMC of the date of first birth and the CMC of the date of birth of the respondent. BASE: All respondents with one or more births (V201 > 0).
V213		Whether the respondent is currently pregnant.
V214		Imputed duration of the current pregnancy. In the imputation process a date of conception of the current pregnancy is calculated from the reported duration of the current pregnancy, if known, or imputed from other available information (see note on imputed dates). The imputed duration of pregnancy is then calculated from that date of conception. BASE: Currently pregnant women (V213 = 1).
V215		Time since last menstrual period as reported by the respondent. The first digit gives the units in which the response was given by the respondent: 1 - Days ago, 2 - Weeks ago, 3 - Months ago, 4 - Years ago, 9 - Special answers. The last two digits give the time since the last period in those units. If the last two digits contain a number greater than 90 then this is a special response. For example, 199 means the response was in days but the number of days was missing on the questionnaire.
V216		Whether the respondent menstruated in the last six weeks is calculated from V215.
V217		Knowledge of the ovulatory cycle indicates when during her monthly cycle the respondent thinks a woman has the greatest chance of becoming pregnant.
V218		Total number of living children is the sum of variables V202 to V205.

<u>Var</u>	<u>Model</u>	<u>Description</u>
V219		Total number of living children including current pregnancy is calculated from V218 by adding 1 if the respondent is pregnant.
V220		Total number of living children including current pregnancy is a grouping of the previous variable, truncating the number to 6 if it was greater than 6.
V221		Interval between the first marriage and first birth in months. If the first birth was prior to the first marriage then this variable is coded 996 "Negative interval." BASE: Ever-married women who have had one or more births ($V501 > 0$ & $V201 > 0$).
V222		Interval between the last birth and the date of the interview in months. BASE: Respondents who have had one or more births ($V201 > 0$).
V223		Completeness of information relating to the date of conception of the current pregnancy. This variable indicates whether the date of conception was exactly specified by the duration of the current pregnancy or the duration was imputed from other information (see note on imputed dates). Codes are different from those used in DHS I.
V224		Number of entries in the birth history (REC21). This variable is also the index to the first birth in the birth history. If there are fewer than twenty births then this is the same as V201 (number of children ever born), but if there are more than twenty births then this will be 20, while V201 gives the full number.
V225		At the time the respondent became pregnant with the current pregnancy, whether the current pregnancy was wanted then, later or not at all. BASE: Currently pregnant women ($V213 = 1$).
V226		Computed time since the last menstrual period. This is computed from the response for V215, with durations exceeding the interval since the last birth ($V227 = 7, 9$) recoded to the response "Before last birth" (code 995) and inconsistent responses flagged on variable V227 (codes 1-6) recoded to 997.
V227		Flag variable indicating inconsistencies found in editing the response for variable V215. 0 No flag 1 Duration given is greater than the interval since the last birth and the respondent did <u>not</u> say, in the maternity section, that she was still amenorrheic since her last birth 2 Duration given plus the duration of amenorrhea after the last birth is greater than the interval since the last birth 3 Duration was reported, but the respondent's period had not returned since the last birth 4 Respondent reported her last period was before her last birth, but she had never given birth 5 Respondent reported never having menstruated, but reported in the maternity section that her period had returned after her last birth 6 Respondent reported her last period was before her last birth, but reported in the maternity section that her period had returned after her last birth 7 Respondent reported a duration since her last period, but this would place her last period <u>during</u> her last pregnancy 8 Respondent reported never having menstruated, but she had children 9 Duration was reported, but the duration would place her period before her last birth

<u>Var</u>	<u>Model</u>	<u>Description</u>
V228	A	Whether the respondent ever had a pregnancy that terminated in a miscarriage, abortion, or still birth, i.e., did not result in a live birth.

Pregnancy terminations

Variables V229 to V234 relate to pregnancy terminations (pregnancies that did not result in a live birth).

BASE: Respondents who have had one or more terminated pregnancies (V228 = 1).

V229	A	Month of the last pregnancy termination.
V230	A	Year of the last pregnancy termination. The date of last termination is taken from a single question in the body of the questionnaire. This variable now occupies 4 digits.
V231	A	Century month code of the last pregnancy termination. The CMC date of termination is calculated from the preceding questions, or from the calendar, if possible, in cases where an exact date was not given for the date of last pregnancy termination.
V232	A	Date flag for the last terminated pregnancy.
V233	A	Months pregnant when the pregnancy terminated. BASE: Respondents who have had one or more terminated pregnancies since the cutoff date for the calendar/health section (V228 = 1 & V231 >= V017).
V234	A	Whether the respondent had other pregnancy terminations before the last one. BASE: Respondents who have had one or more terminated pregnancies since the cutoff date for the calendar/health section (V228 = 1 & V231 >= V017).
V235	A	Birth history index for last child born prior to the start of the calendar. Index is zero if no child was born before the start of the calendar.
V236	—	Whether the interval between the last birth and the date of interview was four calendar years or more. See B14 for more explanation. BASE: Women with at least one birth (V201 > 0).
V237		Whether there were any other live births in the interval between the last birth and the date of interview. See B15 for more explanation.
V238		Total number of births in the last three years is defined as all births in the months 0 to 35 prior to the month of interview, where month 0 is the month of interview.

Pregnancies terminated prior to beginning of calendar

V239		Whether or not, the respondent had pregnancies that miscarried, aborted or ended in still birth before calendar beginning
V240		Month of last termination prior to calendar
V241		Year of last termination prior to calendar
V242		CMC termination ended prior to calendar
V243		Completeness of information for the last termination date prior to calendar

Section 31 (REC31)

Contraceptive Table

Var Model Description

- V301 Knowledge of any method is classified into modern, traditional and folkloric methods as follows: Modern methods are Pill, IUD, Injections, Diaphragm, Condom, Female Sterilization, Male Sterilization, Implants, female condom, Foam/Jelly and lactational amenorrhea. Traditional methods are Periodic Abstinence (Rhythm), Withdrawal, and Abstinence. Folkloric methods are the category "other". If a respondent knows both a traditional method and a modern method then the modern method takes priority and she is coded as knowing a modern method. Similarly, if a woman knows a traditional method and a folkloric method, the traditional method takes priority.
- V302 Ever use of a modern, traditional or folkloric method is created in the same way as V301.

Contraceptive Table

The contraceptive table contains entries for 20 contraceptive methods, and for each entry gives information relating to knowledge of the method, ever use of the method, and knowledge of a source for the method. Entries 1 to 15 are standard but entries 15 to 20 are used for country-specific methods. The methods relating to each entry are as follows:

1	Pill	11	Norplant™ or implants
2	IUD	12	Abstinence
3	Injections	13	Lactational amenorrhea
4	Diaphragm	14	Female condom
5	Condom	15	Foam and jelly
6	Female Sterilization	16	
7	Male Sterilization	17	Country specific method 1
8	Periodic Abstinence (Rhythm)	18	Country specific method 2
9	Withdrawal	19	Country specific method 3
10	Other methods	20	Country specific method 4

For Foam/Jelly, if questions about the methods are asked separately (for example, foaming tablets in one set of questions and jelly combined in another set of questions), the original responses are recorded as country-specific variables and the standard variables presented in this section are a composite of the two sets of questions.

In general, the list of the methods used in the contraceptive table does not exist. The variable that is used to recognize these methods is V312. In some cases additional codes are used in the contraceptive table and these should exist in the documentation of the distributed data file.

The contraceptive table contains variables V304A to V3076 as follows:

- V304A Whether the method is modern, traditional or folkloric.
- V304 Knowledge of the method. If questions relating to the method were not asked in a particular country then code 8 "Not asked" is used.

Var Model Description

- V305 Whether the respondent has ever used the contraceptive method.
 BASE: Respondents who knew of the method according to V304, either spontaneously (1) or after probing (being read a description of the method) (2).
- V307 Method currently used. This is a multiple occurrence variable that gives all the methods used by the respondent.
 BASE: Respondents who ever used the method according to V305.

Example Contraceptive Table:

	1	2	3	4	5	6	7	8	9	10	11	12	13	...	20
	Pill	IUD	Inj	Dia	Cond F.	St.M.	St.P.Ab.	With	Other	Norpl	Abst	CS1	...	CS8	
V304A	1	1	1	1	1	1	1	2	2	3	1	2	3	...	3
V304	1	0	0	0	1	0	0	1	0	0	8	8	8	...	8
V305	0				1	0		1	0						
V307								1							

In this example the entries in the table are shown across the page while the variables in each entry are shown down the page. The numbers shown above the method names are the occurrence or entry number associated with that method. The respondent knew three methods, Pill, Condom and Periodic Abstinence. The respondent has used Condoms and Periodic Abstinence. The respondent is using periodic abstinence.

Section 32 (REC32)

Contraceptive Use

<u>Var</u>	<u>Model</u>	<u>Description</u>
V310		Number of living children at the time the respondent first used a contraceptive method. BASE: All respondents who have ever used a contraceptive method (V302 > 0).
V311		Number of living children at the time of first use is a grouped form of V310, with 4 or more truncated to 4 and respondents who have never used a contraceptive method coded 5.
V312		Current contraceptive method. Pregnant women are coded 0 "Not currently using."
V313		Type of contraceptive method categorizes the current contraceptive method as either a modern method, a traditional method, or a folkloric method.
V314	—————	Method of periodic abstinence indicates how the respondent determined on which days to abstain from sexual intercourse the last time the respondent used periodic abstinence. BASE: Current users of periodic abstinence (V312 = 8) – Note that this is different from the base used in DHS I.
V315		Month started using a contraceptive method by the respondent or her current partner (see note on imputed dates).
V316		Year started using a contraceptive method by the respondent or her current partner (see note on imputed dates). This variable now occupies 4 digits.
V317		Century month code for the date started using a contraceptive method by (see note on century month codes).
V318		Completeness of information for the date of sterilization of the respondent or her partner (see note on imputed dates). Codes for DHS III are different from the codes used in DHS I.

Sterilization.

Variables V319 to V322 relate to dates of sterilization.

BASE: Women who are sterilized or whose partner is sterilized (V312 = 6 or V312 = 7).

V319		Years since sterilization in 2-year groups, truncated at 10 years.
V320		Age at sterilization in 5-year groups <25, 25-29, 30-34, 35-39, 40-44, 45-49.
V321		Marital duration at sterilization in 5-year groups with single women and those sterilized before marriage coded 0.
V322		Parity at sterilization, truncated at 5+ children.

Pill Use.

~~Variables V323 and V325 relate to the use of the pill. See variable V372 to V374 for additional information relating to pill use.~~

~~BASE: Respondents currently using the pill (V312 = 1).~~

V323	—————	Brand of pill currently being used by the respondent. Codes are country specific.
V325	A	Cost of pills in the local currency. The width of this field has been increased to six characters to accommodate a variety of currencies. Code 999996 indicates that the pills were

provided free and code 999998 indicates that the respondent did not know the price of the pills.

Source of modern contraceptive methods.

Variables V326 to V328 and V327 relate to sources of contraception for current users of modern methods. See variables V379 to V380 for additional information relating to the source of modern contraceptive methods.

BASE: Respondents currently using a modern method (V312 >= 1 & V312 <= 7 or V312 = 11 or V312 >= 13 & V312 <= 16).

- V326 The last source visited to obtain the current modern contraceptive method. Codes are country-specific, but the major categories are standard.
- V327 The last source visited for users of modern methods in standard coding groups constructed from V326. The standard coding categories for this variable have been changed to separate non-governmental organizations (NGOs) from other private sector sources.
- ~~V328 Main reason the respondent selected the source for the method, rather than using a different source for the method.~~

Current use of contraception.

Variables V337 relates to the current use of contraception.

BASE: Current users of contraception (V312 <> 0).

- V337 Months of use of the current contraceptive method. For Model "B" questionnaires, if the number of months of use exceeds 8 years, it is coded as 96 For Model "A" questionnaires, if the woman is using the method throughout the calendar, the duration is coded as 95.

Last method discontinued in the last five years.

Variables V359 and V360 relate to the last method discontinued in the last five years. The information for these variables is taken from the calendar.

BASE: Respondents who discontinued use of a method in the last five years.

- V359 A Last method discontinued in the last five years.
- V360 A Reason for the discontinuation of the last method discontinued in the last five years.

Pattern and intentions for future use.

Variables V361 to V364 relate to the respondent's past contraceptive practice and future intentions for using contraception.

- V361 Pattern of past contraceptive use. For model "B" questionnaires the questions relating to contraceptive use since the last birth are not asked and thus the respondent cannot be categorized as having used a method since the last birth or having only used a method before the last birth. In countries using the model "B" questionnaire, all past users are given code 3. In countries using model "A" questionnaires, all women who have not used in the calendar are treated as past users and given code 3.

<u>Var</u>	<u>Model</u>	<u>Description</u>
V362		Intention to use a contraceptive method in the future is based on two questions in the model questionnaires, and classifies those intending to use a method in the future by whether they intend to use that method in the next twelve months or not. The two "Unsure" categories correspond to replies of unsure about using a method in the future (unsure about use) or, for those intending to use a method in the future, unsure about whether they intend to use that method in the next twelve months (unsure about timing). In some countries, women who had never had sexual intercourse were not asked these questions, and are coded 6 on V362. BASE: All respondents not currently using contraception (V312 = 0).
V363		Preferred future method for respondents intending to use a method in the future. BASE: Respondents not currently using a method, but intending to use a method in the future (V312 = 0 & (V362 = 1 or V362 = 2 or V362 = 3)).
V364		Contraceptive use and intention shows current users of modern methods, current users of traditional methods, non-users who intend to use in the future and non-users not intending to use a method. In some countries, women who had never had sexual intercourse were not asked the questions relating to their intention to use contraception in the future, and are coded 5 on V364.
V366A	————	Acceptability of family planning messages being provided on radio.
V366B	————	Acceptability of family planning messages being provided on television. Variables V366A and V366B replace the single variable V366 used in DHS II.
V367		Whether the last child born in the last three/five years was wanted at that time, later or not at all. BASE: Women who gave birth to a child in the last three/five years (V417 > 0).

First contraceptive method used.

~~Variable V369 to V369B relates to the first contraceptive method ever used.~~

V369	A	The first contraceptive method ever used by the respondent. Never users are coded 0.
V369B	B	Whether the first use of contraception was for spacing or limiting reasons, or for other reasons. BASE: Ever users of contraception (V302 <math>\neq</math> 0).

Pill Use.

~~Variable V372 relates to the use of the pill. BASE: Current users of the pill (V312 = 1).~~

V372	————	Whether the package of pills currently being used by the respondent was seen by the interviewer.
-----------------	-----------------	---

Reasons for Non-Use of Contraceptive Methods.

V375A		Reason the respondent is not using a method of contraception to avoid pregnancy.
-------	--	--

Var Model Description

BASE: Women who are not currently using a contraceptive method and who are not pregnant (V312 = 0 & V213 <> 1).

- V376 Reason the respondent does not intend to use a method of contraception in the future. The coding categories have changed in DHS III from those used in DHS II.
BASE: All women not currently using a contraceptive method and not intending to use a method in the future (V362 = 5).
- V376A Whether the respondent would use a method in the future if she was married.
BASE: Women who says they do not intend to use a method because they are not married (V376 = 11).

Sources of Contraception.

V379 Source of any method of contraception is formed from a combination of responses. For current users of modern methods, it is the source of that method. For women who are not currently using any method, it is a source from which they know they can obtain family planning methods, if they know any source. This is not, in general, part of the standard questionnaire since it is replaced with a multiple-choice question. However it is left as a standard variable in case the question asked has only one answer.

V380 Source of any method of contraception coded in standard coding categories is created from V379.

Whether the respondent has heard about family planning in the last few months from any of the following sources:

- V384A On the radio.
V384B On the television.
V384C In a newspaper or magazine.
~~V384D From a poster.~~
~~V384E From leaflets or brochures~~

Family Planning Related Visits

- V393 Whether the respondent was visited by a family planning worker in the twelve months preceding the interview.
- V394 Whether the respondent visited a health facility for any reason in the twelve months preceding the interview.
- V395 Whether any of the staff at the health facility talked to the respondent about family planning.
BASE: Women who visited a health facility in the preceding 12 months (V394 = 1).

Lactational Amenorrhea Method

- ~~V396 Whether the respondent thinks that breastfeeding can affect a woman's chance of becoming pregnant.~~
- ~~V397 Whether the respondent thinks that a woman's chance of becoming pregnant will be increased, decreased, unchanged by breastfeeding or may depend on other factors.~~

Var Model Description

Women responding "No" on variable V396 are coded 0 (Unchanged) on V397.

~~V398~~ ~~Whether the respondent has ever relied on breastfeeding as a method of avoiding pregnancy. BASE: All women who have had at least one birth except those that believe the chance of pregnancy is unchanged or increased by breastfeeding (V201 > 0 & V397 <= 0 & V397 <= 1).~~

~~V399~~ ~~Whether the respondent is currently relying on breastfeeding to avoid getting pregnant. BASE: Respondents who reported ever relying on breastfeeding to avoid pregnancy and who are not currently pregnant and not sterilized (V398 = 1 & V213 <= 1 & V312 <= 6 & V312 <= 7).~~

V3A00A-Z Source of family planning for non users. This question has multiple coding categories and each category is recorded separately in these variables. Most of the categories are standard (VA00A, B, C, D, E, J, K, L, M, N, S, T, U, X). However, room has been left for country-specific categories (VA00F, G, H, I, O, P, Q, R, V, W). Any category not used in a particular country is left blank.

Public Sector

V3A00A Government hospital
V3A00B Government health center
V3A00C FP clinic
V3A00D Mobile clinic
V3A00E Fieldworker
V3A00F Country-Specific public sector
V3A00G Country-Specific public sector
V3A00H Country-Specific public sector
V3A00I Other public sector

Medical Private Sector

V3A00J Private hosp/clin
V3A00K Private pharmacy
V3A00L Private doctor
V3A00M Private mobile clinic
V3A00N Fieldworker
V3A00O Country-Specific medical private sector
V3A00P Country-Specific medical private sector
V3A00Q Country-Specific medical private sector
V3A00R Other medical private sector

Other Private Sector

V3A00S Shop
V3A00T Traditional practitioner
V3A00U Friend, relative
V3A00V Country-Specific other private sector
V3A00W Country-Specific other private sector
V3A00X Other
V3A00Y No source
V3A00Z Any source

V3A01 Before the sterilization operation, whether the respondent was told that sterilization would mean no more children.

<u>Var</u>	<u>Model</u>	<u>Description</u>
V3A02		Told about side effects or problems the respondent might have when first obtained the method she is currently using.
V3A03		Told about side effects by health or FP worker
V3A04		Told how to deal with side effects
V3A05		Told about the availability of other FP methods than the one she is using.
V3A06		Told about other FP methods by health or FP worker
V3A07		First source for current method. This variable is grouped into 4 major categories:
	10	Public sector
	20	Private medical
	30	Other private
	96	Other
V3A08A-Z		Reason for not using. This question has multiple coding categories and each category is recorded separately in these variables. Most of the categories are standard (VA00A-T). However, room has been left for country-specific categories (VA00U-W). Any category not used in a particular country is left blank.
V3A08A		Not married
V3A08B		Not having sex
V3A08C		Infrequent sex
V3A08D		Menopausal/hysterectomy
V3A08E		Subfecund/infecund
V3A08F		Postpartum amenorrheic
V3A08G		Breastfeeding
V3A08H		Fatalistic
V3A08I		Respondent opposed
V3A08J		Husband/partner opposed
V3A08K		Others opposed
V3A08L		Religious prohibition
V3A08M		Knows no method
V3A08N		Knows no source
V3A08O		Health concerns
V3A08P		Fear of side effects
V3A08Q		Lack of access/too far
V3A08R		Costs too much
V3A08S		Inconvenient to use
V3A08T		Interferes with the body's processes
V3A08U		Country specific
V3A08V		Country specific
V3A08W		Country specific
V3A08X		Other
V3A08Z		Don't know

Section 41 (REC41)

Maternity

The maternity history contains up to six entries, relating to births in the three/five years preceding interview. The use of three/five years is country specific. The entries are in reverse order, such that the first entry relates to the last birth in the last three/five years. There is an entry for all children born in the last three/five years including all twins. The period of three/five years includes months 0 to 35/59 prior to the interview, with month 0 being the month of interview. If there are more than six births in the last three/five years then only the last six are included in the maternity history. Each of the following variables, duration of breastfeeding (M5), duration of postpartum amenorrhea (M7) and duration of postpartum abstinence (M9) may have several cases coded 97 "Inconsistent" since the duration of breastfeeding, amenorrhea or abstinence was impossible in the interval between the birth and the following birth or date of interview if the most recent birth (only the date of interview in the case of breastfeeding).

<u>Var</u>	<u>Model</u>	<u>Description</u>
MIDX		Index to the birth history. All births in the last three/five years have entries in this section, and thus the index increases by one each entry. See the example maternity history below. For twins the information in their entries will be identical for all variables relating to prenatal care.
M1		The number of tetanus toxoid injections given during the pregnancy to avoid convulsions after birth. This variable indicated whether the respondent received a tetanus toxoid injection during the pregnancy for DHS-I countries.
M2A-N		The type of person who gave prenatal care to the respondent prior to the birth. This question has multiple coding categories and each category is recorded separately in these variables. Most of the categories are standard (M2A, B, C, F, G, J, N), however room has been left for country-specific categories (M2D, E, H, I, K, L). Any category not used in a particular country is left blank.
M2A		Doctor.
M2B		Nurse/Midwife.
M2C		Auxiliary Midwife.
M2D		Country-specific health professional.
M2E		Country-specific health professional.
M2F	X	Trained (traditional) birth attendant.
M2G		Traditional birth attendant.
M2H	X	Relative. (non-standard and rarely used).
M2I		Country-specific other person.
M2J		Country-specific other person.
M2K		Other responses - uncoded.
M2L		Country-specific other.
M2M		Country-specific other.
M2N		No one.
M3A-N		The type of person who assisted with the delivery of the child. The coding of these variables is the same as for M2A-N, except that the category "Relative" is a standard category (M3H) for this variable.
M4		The duration of breastfeeding of the child in months. The maximum period allowed during the data editing was the interval between the date of birth of the child and the date of interview. Cases which exceeded this duration were left with the original response, but are

<u>Var</u>	<u>Model</u>	<u>Description</u>
		coded with one of the flag codes on variable M27 and were set to code 97 "Inconsistent" on variable M5. The code 96 (breastfed until died) is no longer used.
M5		The calculated months of breastfeeding gives the duration of breastfeeding as in M4, but with the duration calculated if the respondent is still breastfeeding the child or the child was breastfed until it died. Inconsistent durations based on the original reporting of the duration of breastfeeding are recoded to 97. In cases where the duration was one month longer than the interval the duration was shortened by one month, consistent with the "Rule of one" applied in DHS I. If the duration of breastfeeding exceeded the age of death of the child, the duration of breastfeeding was changed to the age at death of the child.
M6		The duration of postpartum amenorrhea after the birth of the child in months. The maximum period allowed during the data editing was the interval between the date of birth of the child and the date of conception of the following child (date birth less nine months was used for the date of conception) or the date of interview if there was no following birth. Cases which exceeded this duration were left with the original response, but are coded with one of the flag codes on variable M28 and were set to code 97 "Inconsistent" on variable M7.
M7		The calculated months of postpartum amenorrhea give the duration of amenorrhea as in M6, but with the duration calculated if the period did not return after the birth and before the following birth or the date of interview. Inconsistent durations based on the original reporting of the duration of amenorrhea are recoded to 97. In cases where the duration was one month longer than the interval the duration was shortened by one month, consistent with the "Rule of one" applied in DHS I.
M8		The duration of postpartum abstinence after the birth of the child in months. The maximum period allowed is calculated in the same way as for M6 and cases exceeding this duration were left with the original response, but are coded with one of the flag codes on variable M29 and are coded 97 "Inconsistent" on M9.
M9		The calculated months of postpartum abstinence give the duration of abstinence as in M8, but with the duration calculated if the respondent was still abstaining after the birth. Inconsistent durations based on the original reporting of the duration of abstinence are recoded to 97. In cases where the duration was one month longer than the interval the duration was shortened by one month, consistent with the "Rule of one" applied in DHS I.
M10		Whether the child was wanted at the time of <u>pregnancy</u> , whether the child was wanted, but later, or whether the child was not wanted at all.
M11		For women who wanted the child later, how much longer the respondent would have preferred to wait. The first digit gives the units in which the respondent gave her answer, code 1 indicates a response in months, code 2 in years, with 9 meaning a special answer was given. The last two digits give the time in the units given. Any value for time greater than 90 is a special answer.
M13		Timing of first antenatal visit for the pregnancy is given in months from the start of the pregnancy. BASE: Women who had seen someone for antenatal care (M2N <> 1).
M14		Number of antenatal visits during the pregnancy. Women who did not see anyone for antenatal care during the pregnancy are coded 0.

Var Model Description

M15	Place of delivery of child. Coding categories are standard and are constructed with a major category for the first digit and a minor category for the second digit. Country-specific codes are added under the respective major coding categories as needed. For example "Home of traditional birth attendant" would be coded 13 since the category relates to a home (major category 1) and categories 11 and 12 are already used as standard categories.
M17	Whether child was born by caesarian section.
M18	Size of child as reported subjectively by the respondent.
M19	Weight of child at birth given in kilograms with three implied decimal places (or grams with no decimal places). Children who were not weighed are coded 9996. In some countries, the birth weight was collected in grams, i.e. a total of four digits, whereas other countries collected the weight in kilograms to one decimal place, i.e. a total of two digits. In the latter case, the third and fourth digit are set to zeros. In a few countries, the weight was collected in pounds and/or ounces. For these countries, the original weight variables are stored as a country-specific variable and this variable contains the weight converted to kilograms.
M19A	Whether the weight at birth (variable M19) was recorded from a health card (code 1) or from the mother's recall (code 2). Children who were not weighed at birth are coded 0.
M21	Reason the respondent stopped breastfeeding the child. Children who breastfed until they died are coded 3 (child died) as the reason stopped breastfeeding. Code 97 indicates cases where data was missing on whether the child was ever breastfed. BASE: Children who are no longer being breastfed, but were ever breastfed (M4 < 94 & M4 < 95).
M27	Flag variable for breastfeeding, indicating types of problems found in editing the duration of breastfeeding.
M28	Flag variable for postpartum amenorrhea, indicating types of problems found in editing the duration of postpartum amenorrhea.
M29	Flag variable for postpartum abstinence, indicating types of problems found in editing the duration of postpartum abstinence. Codes for these flag variables are as follows: 0 No problem. 1 Duration exceeds interval between birth and succeeding birth (for amenorrhea and abstinence) or date of interview (for breastfeeding). 2 Duration exceeds interval but only by one month. In DHS I the duration would have been modified to fit the interval in this case. This modification is made to the calculated months of postpartum amenorrhea (M7), postpartum abstinence (M9) or breastfeeding (M5) starting from DHS III. 3 Duration of breastfeeding exceeds the age at death of the child. 4 Duration of postpartum amenorrhea, postpartum abstinence or breastfeeding extends into the following pregnancy in the calendar. This code only applies to Model A countries.
M30	Whether the respondent had any of the following problems at the time of the birth of the child: Long labor, defined as regular contractions lasting more than 12 hours.

<u>Var</u>	<u>Model</u>	<u>Description</u>
M31	—————	Excessive bleeding that was so much that the respondent feared it was life threatening.
M32	—————	A high fever with a bad smelling vaginal discharge.
M33	—————	Convulsions not caused by fever.
M34		Time after the birth at which the respondent first breastfed the child. The first digit gives the units in which the respondent gave her answer. Code 0 means the child was breastfed immediately after birth, code 1 indicates the response was in hours, code 2 in days, with code 9 meaning a special answer was given. The last two digits give the time in the units given. Any value for time greater than 90 is a special answer. The response "Immediately" is recorded as 000. BASE: Child who were ever breastfed (M4 < 94).
M35		Number of times the child was breastfed during the previous night. BASE: Children still being breastfed (M4 = 95).
M36		Number of times the child was breastfed during the daylight hours the previous day. BASE: Children still being breastfed (M4 = 95).
		Other foods given to the child in the last 24 hours.
M37A		Plain water.
M37B		Sugar water.
M37C		Juice.
M37D		Herbal tea.
M37E		Powdered or tinned milk.
M37F		Baby formula.
M37G		Fresh (cow's or goat's) milk.
M37H-K		<i>Country-specific other liquids.</i>
M37L		Any other liquid.
M37M-O		<i>Country-specific other solid or mushy food.</i>
M37P		Any <u>other</u> solid or semi-solid foods.
M37Q		Food made from wheat, maize, rice, sorghum or other local grains, Examples of such food are porridge, bread or noodles.
M37R		Food made from cassava, plantain, yams or other local tubers.
M37S		Eggs, fish, or poultry.
M37T		Meat.
M37U		Other fruits/vegetables
M37V		Meat, poultry, fish, shellfish, eggs
M37W		Legumes (lentils, beans, peanuts)
M37X		Cheese/yogurt
M37Y		Foods made with oil, fat, butter
M37Z		Country-specific food
M37XX		Country-specific food
M37XY		Country-specific food
M37XZ		Country-specific food BASE: Youngest child living with his/her mother.
M38		Whether the child drank anything from a bottle with a nipple during the previous day and night. BASE: Living children (B5(MIDX) = 1).

<u>Var</u>	<u>Model</u>	<u>Description</u>
M39		The number of times the children received anything to eat, aside from breastmilk, including both meals and snacks. Children who are being exclusively breastfed are coded 0. BASE: Living children (B5(MIDX) = 1).
M40A-XZ		The number of days during the last 7 days on which the child received each of the following liquids and foods:
M40A		Plain water
M40B		Sugar water
M40C		Fruit juice
M40D		Herbal tea
M40E		Powdered or tinned milk
M40F		Commercially produced infant formula
M40G		Fresh milk
M40H		Tinned, powdered or fresh animal milk
M40I		Country specific other liquid
M40J		Country specific other liquid
M40K		Country specific other liquid
M40L		Other liquid
M40M		Pumpkin, carrots, red/yellow yams, red sweet potatoe
M40N		Any green leafy vegetables
M40O		Mango, papaya or other Vitamin A rich fruits
M40P		Other solid, semi-solid
M40Q		Food made from local grain
M40R		Food made from local roots or tuber
M40S		Eggs, fish, poultry
M40T		Meat
M40U		Other fruits/vegetables
M40V		Meat, poultry, fish, shellfish, eggs
M40W		Legumes (lentils, beans, peanuts)
M40X		Cheese/yogurt
M40Y		Foods made with oil, fat, butter
M40Z		Country specific food
M40XX		Country specific food
M40XY		Country specific food
M40XZ		Country specific food
M41		Number of months pregnant at the time of last antenatal visit of last pregnancy

Care given during the last antenatal visit for the pregnancy

BASE: Last births under 36 (60) months

M42A		During pregnancy - weighed
M42B		During pregnancy - height measured
M42C		During pregnancy - blood pressure taken
M42D		During pregnancy - urine sample taken
M42E		During pregnancy - blood sample taken

Pregnancy complications

BASE: Last births under 36 (60) months

M43		Told about pregnancy complications at the time of last antenatal visit
-----	--	--

Var Model Description

M44 Told where to go for pregnancy complications

Iron supplementation

BASE: Last births under 36 (60) months

M45 During pregnancy, given or bought iron tablets/syrup

M46 Days tablets or syrup taken

Vision during pregnancy

BASE: Last births under 36 (60) months

M47 During pregnancy, had difficulty with daylight vision

M48 During pregnancy, had difficulty with night blindness

Medication taken to prevent from Malaria

BASE: Last births under 36 (60) months

M49A During pregnancy - took Fansidar for Malaria

M49B During pregnancy - took Chloroquine for Malaria

M49C During pregnancy - took Unknown Drug for Malaria

M49D During pregnancy - took country specific drug for Malaria

M49E During pregnancy - took country specific drug for Malaria

M49F During pregnancy - took country specific drug for Malaria

M49G During pregnancy - took country specific drug for Malaria

M49X During pregnancy - took other drug for Malaria

M49Z During pregnancy - took no drug for Malaria

Check up after delivery

BASE: Last birth under 36(60) months.

M50 After birth, received a check by a health professional

M51 Checkup after delivery timing

M52 After birth, health professional checked health

M53 Place for checkup

M54 Received Vitamin A dose in first 2 months after delivery

Liquids given before milk began to flow

M55A First 3 days, given milk other than breast milk

M55B First 3 days, given plain water

M55C First 3 days, given sugar/glucose water

M55D First 3 days, given gripe water

M55E First 3 days, given sugar/salt/water solution

M55F First 3 days, given fruit juice

M55G First 3 days, given infant formula

M55H First 3 days, given tea/infusions

M55I First 3 days, given honey

M55J First 3 days, given country specific

M55K First 3 days, given country specific

M55L First 3 days, given country specific

Var Model Description

M55M First 3 days, given country specific
M55N First 3 days, given country specific
M55X First 3 days, given other
M55Z First 3 days, given nothing

M56 Sugar added to any of foods or liquids consumed yesterday.

Example Maternity History:

MIDX	1	2	3
M1	1	—	—
M2A-N	010__0_0_0__0	_____	_____
M3A-N	010__00000__0	010__00000__0	010__00000__0
M4	95	15	22
M5	19	15	22
M6	12	12	20
M7	12	12	20
M8	96	96	12
M9	19	19	12
M10	1	1	1
M11	—	—	—
M13	09	—	—
M14	01	—	—
M15	23	23	24
M17	0	0	0
M18	4	3	2
M19	2020	1720	9998
M19A	1	1	9
M27	0	0	0
M28	0	0	0
M29	0	0	0
M34	000	000	000
M35	02	—	—
M36	03	—	—
M37A-XZ	0_0_0_1__0010_10__00010__	_____	_____
M38	0	—	0
M39	5	—	5
M40A-XZ	5_0_0_7__0050_40__03045__	_____	_____
M41	09	—	—
M42A-E	11110	_____	_____
M43	0	—	—
M44	—	—	—
M45	1	—	—
M46	010	_____	_____
M47	0	—	—
M48	0	—	—
M49A-Z	0100__00	_____	_____
M50	—	—	—
M51	—	—	—
M52	—	—	—
M53	—	—	—
M54	1	—	—
M55A	000000000__01	000000000__01	000000000__01
M56	0	—	1

In this example, based on the birth history example, there are three entries representing the three children born in the last three/five years. The first two entries relate to twins and so most of their prenatal care information are identical. The respondent received a tetanus injection, prenatal care from a trained nurse, and delivery assistance from nurse/midwife at a government health post, with the assistance of a relative. One of the twins was still being breastfed, 19 months after the birth, while the other had stopped breastfeeding after 15 months. The respondent's periods had returned after the last birth and the respondent is still

abstaining from sexual relations for 19 months after the birth. Both the twins were wanted at that point in time. The respondent received antenatal care during her pregnancy. The first visit was made after nine months, and she made one visit in total. The twins were both measured at birth and weighed 2.020 kilos and 1.720 kilos, respectively. For child 3 no tetanus injection was given, no prenatal care and the child was delivered with the assistance of a nurse/midwife. The child was breastfed for 22 months, the reported duration of postpartum amenorrhea was of 20 months, and the respondent abstained from sexual relations for 12 months after the birth of this child. The child was larger than average at birth, but had not been weighed at birth. The child was breastfed for 22 months. None of the children of the children were born by caesarian section. The last child living with his mother had been given supplemental foods. This child received fresh milk last night, he received plain water on five days during the last week.

Section 42 (REC42)

Maternity and Feeding

<u>Var</u>	<u>Model</u>	<u>Description</u>
V401	A	Whether the last child born in the last three/five years was born by caesarean section. BASE: Respondents who have had one or more births in the three/five years preceding the survey (V417 > 0).
V404		Whether the respondent is currently breastfeeding a child. This is based on the entries in the maternity history for children born in the last three/five years. If no child was born in the last three/five years, the respondent is assumed not to be breastfeeding. This variable is created by looking for any child which is still being breastfed, and not just whether the last child is being breastfed.
V405		Whether the respondent is currently postpartum amenorrheic. This variable is created from the maternity history by checking if the period returned after the last birth. If the woman is currently pregnant then she is coded as not currently amenorrheic, irrespective of whether her period returned after the last birth. If there are no births in the last three/five years then this variable is coded 0 "Not currently amenorrheic."
V406		Whether the respondent is currently postpartum abstaining. This variable is created from the maternity history by checking if the respondent has resumed sexual relations since the last birth. If there are no births in the last three/five years then this variable is coded 0 "Not currently abstaining."
V407		Number of times the last child was breastfed during the previous night. BASE: Respondents still breastfeeding the last child (V404 = 1).
V408		Number of times the last child was breastfed during the daylight hours the previous day. BASE: Respondents still breastfeeding the last child (V404 = 1).
<u>Other foods given to the child in the last 24 hours:</u>		
V409		Plain water.
V409A		Sugar water.
V410		Juice.
V410A		Herbal tea.
V411		Powdered or tinned milk.
V411A		Baby formula.
V412		Fresh (cow's or goat's) milk.
V413A-D		Country-specific other liquids.
V413		Any other liquid.
V414A-C		Country-specific other solid or mushy food.
V414D		Any other solid or semi-solid food.
V414E		Food made from wheat, maize, rice, sorghum or other local grains, Examples of such food are porridge, bread or noodles.
V414F		Food made from cassava, plantain, yams or other local tubers.
V414G		Eggs, fish, or poultry.
V414H		Meat.
		BASE: Last born child is still alive (B5(1) = 1).
V415		Whether the child drank anything from a bottle with a nipple the previous day and night. BASE: Respondents whose last child born in the last three/five years was still alive (V417 > 0 & B5(1) = 1).

Var Model Description

V416	Whether the respondent has heard of the special-named oral rehydration product for treating children with diarrhea. This variable is coded 1 if the respondent had used the ORS product to treat a child in the previous two weeks, 2 if the respondent had heard of the ORS product. BASE: All respondents.
V417	Number of entries in the maternity history.
V418	Number of entries in the health history.
V419	Number of entries in the height and weight table.
V420	Code assigned to the person measuring the children for the height and weight section. Codes are country-specific. This variable now occupies 3 characters. BASE: Respondents who had given birth to a child in the last three/five years (V417 > 0).
V421	Code assigned to the assistant measurer. Codes are country-specific. . This variable now occupies 3 characters. BASE: Respondents who had given birth to a child in the last three/five years (V417 > 0).
V426	Time after the birth at which the respondent first breastfed the last child. The first digit gives the units in which the respondent gave her answer. Code 0 means the child was breastfed immediately after birth, code 1 indicates the response was in hours, code 2 in days, with code 9 meaning a special answer was given. The last two digits give the time in the units given. Any value for time greater than 90 is a special answer. The response "Immediately" is recorded as 000. BASE: Respondents whose last child born in the last three/five years was ever breastfed. (V208V417 > 0 & M4(1) <> 94).

Maternal Anthropometry

Data on maternal anthropometry is only collected for mothers of children born in the three/five years preceding the survey (months 0 to 59 before the survey).
BASE: Mothers of children born in the preceding three/five years (V417 > 0).

V436	X Upper arm circumference of the respondent in centimeters. There is one implied decimal place in the arm circumference (decimal points are not included in the data file). To produce the arm circumference in centimeters divide by 10.
V437	Weight of the respondent in kilograms. There is one implied decimal place in the weight (decimal points are not included in the data file). To produce the weight in kilograms divide by 10.
V438	Height of the respondent in centimeters. There is one implied decimal place in the height (decimal points are not included in the data file). To produce the height in centimeters divide by 10.
V439	Height for Age percentile.
V440	Height for Age standard deviations from the reference median.
V441	Height for Age percent of reference median.
V442	Weight for Height percent of reference median based on DHS reference standard.

<u>Var</u>	<u>Model</u>	<u>Description</u>
V443		Weight for Height percent of reference median based on Metropolitan Life or Fogarty reference standard.
V444		Weight for Height percent of reference median based on WHO reference standard. The NCHS/FELS/CDC reference standard only contains data for children up to the age of 18 years. For all women aged 18 and over, the value of 215 months (17 years, 11 months) is used for their age, on the assumption that women are fully grown by the age of 18. Weight-for-age indices are not included as the weight of an adult woman is very dependent on her height. For the weight-for-height indicators, the CDC standard only applies up to a height of 137 centimeters, and almost all adult women are taller than this height. For this reason the weight-for-height Z-scores and percentiles are not available. However, three measures of percent of reference median are included, one based on the Metropolitan Life or Fogarty standard, the second based on the WHO standard and a third based on a DHS standard. These indices have been adjusted for pregnant women according to duration of pregnancy. The anthropometric indices above are based on the CDC Standard Deviation-derived Growth Reference Curves derived from the NCHS/FELS/CDC Reference Population. The measures are presented with two implied decimal places (no decimal points are included in the data file). To produce the actual measure, divide the variable by 100. If either the weight or the height of the respondent is missing, then the corresponding measures above are set to the missing code 9999 or 99999. If either the height or the weight is outside of the acceptable range for the calculation of these measures, then the corresponding measures are set to code 9998 or 99998.
V444A		Weight for Height standard deviations from the reference median based on the DHS reference standard.
V445		Body mass index (BMI), or Quetelet's index, for the respondent is defined as her weight in kilograms divided by the square of her height in meters (W/H^2). There are two implied decimal place in the BMI (decimal points are not included in the data file). To produce the BMI divide by 100. The BMI has <u>not</u> been adjusted for pregnant women.
V446		Rohrer's index for the respondent is defined as her weight in kilograms divided by her height in meters cubed (W/H^3). There are two implied decimal place in the Rohrer's index (decimal points are not included in the data file). To produce the Rohrer's index divide by 100. The Rohrer's index has <u>not</u> been adjusted for pregnant women.
V447		Result of measurement of the respondent. Fully measured women are coded 0, and reasons for not measuring the respondent are coded 2 and above (see HW13).
V447A		Women's age in years from household report
V448		Drinking pattern with diarrhea indicates what the respondent believes a child should be given to drink when the child has diarrhea.
V449		Eating pattern with diarrhea indicates what the respondent believes a child should be given to eat when the child has diarrhea.
V450A M,X,Z		The signs of illness with diarrhea that would indicate to the respondent that the child should be taken to a health facility or health worker for treatment.
V450A		Repeated watery stools.

Var Model Description

V450B — Any watery stools.
 V450C — Repeated vomiting.
 V450D — Any vomiting.
 V450E — Blood in the stools.
 V450F — Fever.
 V450G — Marked thirst.
 V450H — Not eating well or not drinking well.
 V450I — Getting sicker or very sick.
 V450J — Not getting better.
 V450K — *Country specific.*
 V450L — *Country specific.*
 V450M — *Country specific.*
 V450X — Other responses.
 V450Z — Does not know any signs of illness with diarrhea.

~~V451A-K,X,Z — The signs of illness with a cough that would indicate to the respondent that a child should be taken to a health facility or a health worker for treatment.~~

V451A — Fast breathing.
 V451B — Difficult breathing.
 V451C — Noisy breathing.
 V451D — Fever.
 V451E — Unable to drink.
 V451F — Not eating well or not drinking well.
 V451G — Getting sicker or very sick.
 V451H — Not getting better.
 V451I — *Country specific.*
 V451J — *Country specific.*
 V451K — *Country specific.*
 V451X — Other responses.

Respondents 15-17

V452A Under age 18 (HH report). Respondents aged 15-17 have their caretaker line number collected and used for getting the consent to draw blood for anemia testing.
 V452B Line number of parent/responsible adult

Anemia testing

V452C Read consent statement. A consent statement is read to the respondent for agreement before taking a sample of blood.
 V453 Hemoglobin level (g/dl - 1 decimal)
 V454 Currently pregnant (household report)
 V455 Result of measuring (Hemoglobin)
 V456 Hemoglobin level adjusted by altitude (g/dl - 1 decimal)
 V457 Anemia level
 V458 Agrees to referral. In case where the anemia level is severe, respondents are asked whether this information can be given to a doctor at a specified health facility for follow up.

Var Model Description

Bednets in the household

V459 Have bednet for sleeping (household report)
V460 Children under 5 slept under bednet last night (HH report)
V461 Respondent slept under bednet

V462 Washed hands before preparing last meal

Tobacco use

Type of tobacco the respondent currently smokes

V463A Cigarettes
V463B Pipe
V463C Other tobacco
V463D Country specific
V463E Country specific
V463F Country specific
V463G Country specific
V463Z Does not smoke

V464 Number of cigarettes in last 24 hours
BASE: Respondents smoking cigarettes.

V465 Disposal of youngest child's stools when not using toilet.
BASE: Respondents with children under 5 living with them.

V466 When child is seriously ill, the respondent can decide by herself whether or not the child should be taken for medical treatment.
BASE: Respondents with children living with them.

Getting medical care for herself

In case where the respondent is sick, this set of questions give an answer to the major problems preventing her from getting a medical advice or treatment.

V467A Knowing where to go
V467B Getting permission to go
V467C Getting money needed for treatment
V467D Distance to the health facility
V467E Having to take transport
V467F Not wanting to go alone
V467G Concern that there may not be a female health provider

V468 Column used for variables related only to last births. In the standard, some questions are asked about the last child but in some countries these questions are asked about all children born in the last three/five years. This variable gives an indication on whether the questionnaire followed the standard or was changed to include more children.

Times last under three/five child living with his mother was given supplements during last 24 hours

Var Model Description

V469A Plain water
V469B Sugar water
V469C Fruit juice
V469D Herbal tea
V469E Powdered/tinned milk
V469F Commercially produced baby formula
V469G Fresh milk
V469H Tinned, powdered or fresh animal milk
V469I Country specific other liquid
V469J Country specific other liquid
V469K Country specific other liquid
V469L Other liquid
V469M Pumpkin, carrots, red/yellow yams, red sweet potato
V469N Any green leafy vegetables
V469O Mango, papaya or other Vitamin A rich fruits
V469P Other solid, semi-solid foods
V469Q Food made from local grain
V469R Food made from local roots/tuber
V469S Eggs, fish, poultry
V469T Meat
V469U Other fruits/vegetables
V469V Meat, poultry, fish, shellfish, eggs
V469W Legumes (lentils, beans, peanuts)
V469X Cheese/yogurt
V469Y Foods made with oil, fat, butter
V469Z Country specific food
V469XX Country specific food
V469XY Country specific food
V469XZ Country specific food

Times last under three/five child living with his mother was given supplements during last 7 days

V470A Plain water
V470B Sugar water
V470C Fruit juice
V470D Herbal tea
V470E Powdered/tinned milk
V470F Commercially produced baby formula
V470G Fresh milk
V470H Tinned, powdered or fresh animal milk
V470I Country specific other liquid
V470J Country specific other liquid
V470K Country specific other liquid
V470L Other liquid
V470M Pumpkin, carrots, red/yellow yams, red sweet potato
V470N Any green leafy vegetables
V470O Mango, papaya or other Vitamin A rich fruits
V470P Other solid, semi-solid foods
V470Q Food made from local grain

<u>Var</u>	<u>Model</u>	<u>Description</u>
V470R		Food made from local roots/tuber
V470S		Eggs, fish, poultry
V470T		Meat
V470U		Other fruits/vegetables
V470V		Meat, poultry, fish, shellfish, eggs
V470W		Legumes (lentils, beans, peanuts)
V470X		Cheese/yogurt
V470Y		Foods made with oil, fat, butter
V470Z		Country specific food
V470XX		Country specific food
V470XY		Country specific food
V470XZ		Country specific food

Section 43 (REC43)

Health History

The health history contains up to six entries, relating to children born in the last three/five years. All children born in the last three/five years, covering months 0 to 35/59 prior to the interview, as for the maternity history, are included. The children who have died are included in this section, whereas in the DHS I individual recode only living children were included. For children who have died, none of the variables are applicable, but the entry is included to facilitate linking with children's data in other sections. If there are more than six children born in the last three/five years then only the last six are included in the health history. See the example health history below.

<u>Var</u>	<u>Model</u>	<u>Description</u>
------------	--------------	--------------------

HIDX		Index to the birth history. <u>All</u> children born in the last three/five years have entries in this section. Children of multiple births each have their own entry as in the maternity history.
------	--	--

H1		Whether the respondent has a health card for the child and whether she could produce it for the interviewer. Code 1 means the interviewer saw the health card for the child, whereas code 2 means the respondent reported she had a health card for the child but the interviewer did not see it. Code 3 indicates that the respondent had a health card for the child at some point in time, but no longer has the health card. The health card is used to verify whether specific vaccinations were given and to record the dates of vaccination of the children rather than asking the respondent to report vaccinations.
----	--	--

H2		Whether a date of vaccination was recorded on the health card for BCG. Code 1 means the child has a date recorded for the vaccination. Code 2 is used to indicate that the respondent reported that the child had received the vaccination although the health card was not seen or did not exist, or the vaccination was not recorded on the health card, but was reported by the mother. Code 3 is used to indicate situations where the health card is clearly marked to indicate that the vaccination was given, but no date was recorded on the health card for the vaccination.
----	--	---

H2D		BCG vaccination date - day.
-----	--	-----------------------------

H2M		BCG vaccination date - month.
-----	--	-------------------------------

H2Y		BCG vaccination date - year. This variable now occupies 4 digits.
-----	--	---

H3		DPT 1 vaccination. As for H2, H2D, H2M, H2Y.
----	--	--

H4		Polio 1 vaccination. As for H2, H2D, H2M, H2Y.
----	--	--

H5		DPT 2 vaccination. As for H2, H2D, H2M, H2Y.
----	--	--

H6		Polio 2 vaccination. As for H2, H2D, H2M, H2Y.
----	--	--

H7		DPT 3 vaccination. As for H2, H2D, H2M, H2Y.
----	--	--

H8		Polio 3 vaccination. As for H2, H2D, H2M, H2Y.
----	--	--

H9		Measles vaccination. As for H2, H2D, H2M, H2Y.
----	--	--

H0		Polio 0 (at birth) As for H2, H2D, H2M, H2Y.
----	--	--

If the vaccination date reported is inconsistent with the date of birth or the date of interview or with the dates of other vaccinations part or all of the date of vaccination may be set to 97 "Inconsistent."

BASE: Children who have the vaccination recorded on the health card (H2 = 1).

Var Model Description

H10 Whether the child ever received any vaccination to prevent him/her from getting diseases. This variable comes from a single question in the model questionnaires, which is used if the respondent does not have a health card for the child, and is not a summary of the preceding variables.
 BASE: Children whose mother could not produce a health card (H1 = 0 or H1 = 2 or H1 = 3).

Diarrhea

Variables H11 to H21 relate to the prevalence and treatment of diarrhea.
 BASE: All living children born in the last three/five years for H11, and children having an episode of diarrhea in the last two weeks (H11 = 1 or H11 = 2) for H11A to H21.

H11 Whether the child had diarrhea in the last 24 hours or within the last two weeks. *Code 1 indicates that the child had been ill in the last 24 hours*, code 2 indicates that the child had been ill with diarrhea in the last two weeks. ~~Code 1 is a country-specific code for DHS-III.~~ In case the question about diarrhea in the last 24 hours is used, the code 1 can be used, in this case the code 2 is used to indicate that the child had diarrhea the last two weeks but not in the last 24 hours otherwise the code 2 is for the last 2 weeks including the last 24 hours.

~~H11B Whether there was any blood in the stools during the last episode of diarrhea.
 BASE: Children having an episode of diarrhea in the last two weeks (H11 = 1 or H11 = 2).~~

~~H11C The number of bowel movements on the worst day of diarrhea.
 BASE: Children having an episode of diarrhea in the last two weeks (H11 = 1 or H11 = 2).~~

H12A-X The place at which medical treatment or advice was sought for the last episode of diarrhea. This question has multiple coding categories and each category is recorded separately in these variables. Most of the categories are standard (H12A, B, C, D, E, J, K, L, M, N, S, T, X). However, room has been left for country-specific categories (H12F, G, H, I, O, P, Q, R, U, V, W). Any category not used in a particular country is left blank.

Public Sector.

- H12A Government Hospital.
- H12B Government Health Center.
- H12C Government Health Post.
- H12D Mobile Clinic.
- H12E Community Health Worker.
- H12F *Country-specific public sector.*
- H12G *Country-specific public sector.*
- H12H *Country-specific public sector.*
- H12I Other public sector.

Medical Private Sector.

- H12J Private Hospital or Clinic.
- H12K Pharmacy.
- H12L Private Doctor.
- H12M Mobile Clinic.
- H12N Community Health Worker.
- H12O *Country-specific medical private sector.*
- H12P *Country-specific medical private sector.*
- H12Q *Country-specific medical private sector.*

<u>Var</u>	<u>Model</u>	<u>Description</u>
H12R		Other medical private sector. Other Private Sector.
H12S		Shop.
H12T		Traditional Practitioner.
H12U		<i>Country-specific other private sector.</i>
H12V		<i>Country-specific other private sector.</i>
H12W		<i>Country-specific other private sector.</i>
H12X		Other.
H12Y		Whether no treatment or advice was sought for the diarrhea as reported by the respondent.
H12Z		Whether the child was taken to a medical facility for treatment of the diarrhea. This usually includes being taken to all Public Sector facilities and all Medical Private Sector facilities except for Pharmacy. This variable is a summary of these preceding variables as is used in the final reports.
H13		Whether the child received a sugar-salt-water solution from a special packet (ORS). H13 is coded 1 if the respondent spontaneously reported giving ORS to treat the diarrhea, and code 2 if it was reported only after probing. Code 1 is country specific for DHS III.
H14		Whether the child was given the recommended home solution. H14 is coded 1 if the respondent spontaneously reported giving the recommended home solution to treat the diarrhea, and code 2 if it was reported only after probing. Code 1 is country specific for DHS III and MEASURE DHS+.
H14A		Duration in days for which the child was given a home-made fluid made from recommended ingredients when he/she had diarrhea. BASE: Children receiving a recommended home solution (H14 = 1 or H14 = 2).
H15		Whether the child was given other pills or syrups.
H15A	X	Whether the child was given antibiotics.
H15B		Whether the child was given an injection.
H15C		Whether the child was given an IV (Intravenous feeding).
H15D		Whether the child was given home remedies or herbal medicines.
H15E-H		Country-specific other treatments.
H16		Whether the child was given an increase, the same amount, or a decrease in fluids.
H18	X	Whether the child was given an increase, the same amount, or a decrease in foods.
H20		Whether the child received any other treatment.
H21A		Whether the respondent reported that the child received no treatment.
H21		Whether the child received any treatment or whether advice or treatment was sought for the child. This is a summary of the preceding variables H12A to H12Z, H13, H14, H15 to H15H and H20.

Fever

H22 Whether the child had fever in the last two weeks.

Variables H31 to H40 relate to the prevalence and treatment of cough in the two weeks preceding the interview.

Var Model Description

BASE: All living children born in the last three/five years for H31, and children suffering from cough in the last two weeks for H32 to H38 (H31 = 1 or H31 = 2).

H31	Whether the child had suffered from a cough in the last two weeks <i>and whether the child had been ill with the cough in the last 24 hours</i> . Code 1 indicates that the child had been ill in the last 24 hours, code 2 indicates that the child had been ill with the cough in the last two weeks. Code 1 is country specific for DHS III. In case code 1 is used, code 2 indicates that the child had cough in last two weeks but not in the last 24 hours.
H31B	Whether the child had suffered from rapid breathing when he/she had the cough. BASE: Child who had suffered from a cough (H31 = 1 or H31 = 2).
H32A-X	The place at which medical treatment or advice was sought for the last episode of fever and/or cough. This question has multiple coding categories and each category is recorded separately in these variables. Most of the categories are standard (H32A, B, C, D, E, J, K, L, M, N, S, T, X). However, room has been left for country-specific categories (H32F, G, H, I, O, P, Q, R, U, V, W). Any category not used in a particular country is left blank.
	Public Sector.
H32A	Government Hospital.
H32B	Government Health Center.
H32C	Government Health Post.
H32D	Mobile Clinic.
H32E	Community Health Worker.
H32F	<i>Country-specific public sector.</i>
H32G	<i>Country-specific public sector.</i>
H32H	<i>Country-specific public sector.</i>
H32I	Other public sector.
	Medical Private Sector.
H32J	Private Hospital or Clinic.
H32K	Pharmacy.
H32L	Private Doctor.
H32M	Mobile Clinic.
H32N	Community Health Worker.
H32O	<i>Country-specific medical private sector.</i>
H32P	<i>Country-specific medical private sector.</i>
H32Q	<i>Country-specific medical private sector.</i>
H32R	Other medical private sector.
	Other Private Sector.
H32S	Shop.
H32T	Traditional Practitioner.
H32U	<i>Country-specific other private sector.</i>
H32V	<i>Country-specific other private sector.</i>
H32W	<i>Country-specific other private sector.</i>
H32X	Other.
H32Y	Whether no treatment or advice was sought for the fever and/or cough as reported by the respondent.
H32Z	Whether the child was taken to a medical facility for treatment of the fever and/or cough. This usually includes being taken to all Public Sector facilities and all Medical Private Sector

Var Model Description

facilities except for Pharmacy. This variable is a summary of these preceding variables as is used in the final reports.

Vitamin A vaccination

H33 Received Vitamin A. As for H2.
H33D Vitamin A Day. As for H2D.
H33M Vitamin A month. As for H2M.
H33Y Vitamin A year. As for H2Y.

H34 Whether the respondent received or not a vitamin A dose in form of an ampule, a capsule or syrup in last 6 months

Vaccination campaigns

H35 Any vaccinations in the last 2 years given as part of a national immunization day campaign

H36A Vaccinated during Campaign A
H36B Vaccinated during Campaign B
H36C Vaccinated during Campaign C
H36D Vaccinated during Campaign D
H36E Vaccinated during Campaign E
H36F Vaccinated during Campaign F
BASE for H36A-F: Vaccination campaigns conducted in the area (H35 = 1).

Treatment taken for fever

H37A Fansidar
H37B Chloroquine
H37C Aspirin
H37D Ibuprofen/acetaminophen
H37E Country specific
H37F Country specific
H37G Country specific
H37H Country specific
H37X Other
H37Y Nothing
H37Z Don't know if or what was taken
BASE for H37A-Z: Children under three/five with fever last two weeks (H22 = 1).

Drinking and eating pattern during diarrhea

H38 Amount offered to drink
H39 Amount offered to eat

BASE: Diarrhea last two weeks.

Example Health History:

H1DX	1	2	3
H1	1	—	2
H2	1 22 02 2000	— — — —	2 — — — —

H3	1 24 05 2000	- - - -	2 - - - -
H4	1 24 05 2000	- - - -	2 - - - -
H5	1 29 06 2000	- - - -	2 - - - -
H6	1 29 06 2000	- - - -	2 - - - -
H7	1 25 08 2000	- - - -	2 - - - -
H8	1 25 08 2000	- - - -	2 - - - -
H9	1 30 11 2000	- - - -	8 - - - -
H0	1 22 02 2000	- - - -	0 - - - -
H10	-	-	1
H11	2	-	0
H12A-Z	00__000_0000_000_00000_010	_____	_____
H13	2	-	-
H14	0	-	-
H15	0	-	-
H15A-H	_001_____	_____	_____
H20	0	-	-
H21A	0	-	-
H21	1	-	-
H22	1	-	0
H31	0	-	0
H31B	-	-	-
H32A-Z	00__000_0000_000_00000_010	_____	_____
H33	1 25 _8 2000	- - - -	8 - - - -
H34	0	-	8
H35	1	-	1
H36A-F	0001__	_____	1111__
H37A-Z	011_000_000	_____	_____
H38	5	-	-
H39	3	-	-

In this example, based on the birth history example, there are three entries representing the two living children born in the last three/five years and one child who died. The first two entries are twins, however the second twin has died. The first child has a health card and has dates reported for BCG, DPT 1 & 2 & 3, and Polio 0, 1 & 2 & 3 vaccinations as well as a Vitamin A vaccination.. No questions about Immunization, Health and Nutrition were asked for the dead daughter. The other living child has a health card that the respondent could not produce; the child had received all the vaccinations according to the respondent -- BCG, Polio and DPT with the exception of the Measles for which she does not know whether the child received it or not and the Polio 0 for which she said that the child did not receive it. In the two weeks prior to the interview, the first child listed had an episode of diarrhea but did not see anyone; the mother treated the child by increasing the quantity of fluids the child was given, but and somewhat less to eat. The first child had fever in the two weeks prior to the interview, but was not treated for it in H32A-Z but received Chloroquine and Aspirin in H37A-Z. The first child had received a dose of vitamin A. He received a vaccination during the fourth campaign.

Section 44 (REC44)

Height and Weight

The height and weight table contains information relating to children born in the period three/five years, i.e., 0 to 35/59 months prior to the interview. The entries are in reverse order, such that the youngest child is reported first. All live births in the period are included in the table, including children who have subsequently died.

The information included in this section comes from data collected in the household questionnaire and transferred to the woman's level.

BASE: Living children for variables HW1-HW12, HW14-HW58 (B5(HWIDX) = 1).

<u>Var</u>	<u>Model</u>	<u>Description</u>
HWIDX		Index to the birth history. All live births born in the specified period have entries in this section.
HW1		Age in months of the child is calculated from the country month code of the date of interview less the century month code of the date of birth of the child.
HW2		Weight in kilograms. There is one implied decimal place in the weight (decimal points are not included in the data file). To produce the weight in kilograms divide by 10.
HW3		Height in centimeters. There is one implied decimal place in the height (decimal points are not included in the data file). To produce the height in centimeters divide by 10. Height is supposed to be the recumbent length for children less than 24 months old and the standing height for children born 24 or more months prior to the interview.
HW4		Height for Age percentile.
HW5		Height for Age standard deviations from the reference median.
HW6		Height for Age percent of reference median.
HW7		Weight for Age percentile.
HW8		Weight for Age standard deviations from the reference median.
HW9		Weight for Age percent of reference median.
HW10		Weight for Height percentile.
HW11		Weight for Height standard deviations from the reference median.
HW12		Weight for Height percent of reference median.
		The measures above were calculated using the CDC Standard Deviation-derived Growth Reference Curves derived from the NCHS/FELS/CDC Reference Population. The measures are presented with two implied decimal places (no decimal points are included in the data file). To produce the actual measure, divide the variable by 100. If either the weight or the height of the child is missing then all of the above measures are set to the missing code 9999 or 99999. If either the height or the weight is outside of the acceptable range for the calculation of these measures then all of the above measures is set to code 9998 or 99998.
HW13		Reason the child was not measured. Fully measured children are coded 0, dead children are coded 1, and other reasons for not measuring the child are coded 2 and above.

- HW14 ~~Whether the child has a BCG scar on his/her left shoulder (or any other location used for the BCG vaccination in a particular country).
BASE: Living children (B5(HWIDX)=1).~~
- HW15 Whether the child was measured lying down or standing up. In DHS surveys, children aged less than 24 months are to be measured lying down, children age 24 months or older are to be measured standing up. There may, however, be a considerable discrepancy between policy and practice!
- HW16 Day of birth of the child. This is used in conjunction with the date of measurement of the child to more finely calculate the age of the child in days when computing the anthropometric measures. This level of accuracy can be important in very young children. The age of the child calculated from the day of birth information is not used as a background characteristic for tabulations, but purely to produce a more accurate set of anthropometric indices.
- HW17 Day of measurement.
- HW18 Month of measurement.
- HW19 Year of measurement. This variable now occupies 4 digits.
- ~~HW20 X Upper arm circumference in centimeters. There is one implied decimal place in the upper arm circumference (decimal points are not included in the data file). To produce the upper arm circumference in centimeters divide by 10.~~
- ~~HW21 X Upper arm circumference for Age percentile.~~
- ~~HW22 X Upper arm circumference for Age standard deviations from the reference median.~~
- ~~HW23 X Upper arm circumference for Age percent of reference median.~~
- ~~HW24 X Upper arm circumference for Height percentile.~~
- ~~HW25 X Upper arm circumference for Height standard deviations from the reference median.~~
- ~~HW26 X Upper arm circumference for Height percent of reference median.~~
- HW51 Line number of parent/caretaker
- HW52 Whether the consent to draw a droplet of blood was granted after reading a consent statement to a parent/responsible adult.
- HW53 Hemoglobin level in g/dl with 1 implied decimal
- HW55 Result of measuring (Hemoglobin)
- HW56 Hemoglobin level adjusted by altitude in g/dl with 1 implied decimal
- HW57 Anemia level. Levels below 7.0 g/dl are considered as severe anemia, levels between 7.1g/dl and 9.9g/dl are considered as moderate anemia and cases between 10.0 g/dl and 10.9 g/dl are considered as mild anemia.
- HW58 Agrees to referral. In case where the anemia level is severe, respondents are asked whether this information can be given to a doctor at a specified health facility for follow up

Example Height and Weight Table:

HWIDX	1	2	3
HW1	19	—	47
HW2	90	—	133

<u>Var</u>	<u>Model</u>	<u>Description</u>		
HW3	755		_____	885
HW4	73		_____	9999
HW5	-244		_____	9999
HW6	9093		_____	99999
HW7	151		_____	9999
HW8	-217		_____	9999
HW9	7762		_____	99999
HW10	1241		_____	6909
HW11	-115		_____	50
HW12	9063		_____	10560
HW13	0		_____ 1	0
HW15	1		_____ -	2
HW16	16		_____ -	98
HW17-19	6/9/2001		____/____/____	6/9/2001
HW51	—		_____ -	—
HW52	—		_____ -	—
HW53	—		_____ -	—
HW55	—		_____ -	—
HW56	—		_____ -	—
HW57	—		_____ -	—
HW58	—		_____ -	—

In this example, based on the birth history example, three children are included. The first child aged 19 months, measuring 9.0 kg and 75.5cm. His anthropometric measures are given to 1 implied decimal place. He is measured lying down. The second twin had died (as recorded in HW13). The third child was measured and weighs 13.3 kg and measures 88.5 cm No hemoglobin measurements for both living children.

Section 51 (REC51)

Marriage

<u>Var</u>	<u>Model</u>	<u>Description</u>
V501		Current marital status of the respondent.
V502		Whether the respondent is currently, formerly or never married (or lived with a partner). Currently married includes married women and women living with a partner, and formerly married includes widowed, divorced, separated women and women who have lived with a partner but are not now living with a partner.
V503		Whether the respondent has been married or lived with a man once or more than once. BASE: Ever-married women (V501 > 0).
V504		Whether the partner lives in the household or is now living elsewhere. BASE: Currently married or in union women (V502 = 1).
V505	B	Whether the respondent is in a polygynous union and the number of other wives the respondent's partner currently has. BASE: Currently married or in union women (V502 = 1).
V506	B	The rank of the respondent among the partner's wives. BASE: Currently married or in union women in a polygynous union (V502 = 1 & V505 > 0).

First marriage or union

Variables V507 to V513 relate to the date of start of the first marriage or union.
BASE: Ever-married women (V501 > 0).

V507		Month of start of first marriage or union (see note on imputed dates).
V508		Year of start of first marriage or union (see note on imputed dates). This variable now occupies 4 digits.
V509		Century month code of the date of start of first marriage or union (see note on century month codes).
V510		Completeness of information for the date of start of the first marriage or union (see note on imputed dates). Codes are different from the codes used in DHS I.
V511		Age at start of first marriage or union is calculated from the century month code of the date of start of first marriage or union and the century month code of the date of birth of the respondent.
V512		Years since start of first marriage or union is calculated from the century month code of the start of first marriage or union and the century month code of the date of interview.
V513		Marital duration is actually the number of years elapsed since the start of the first marriage or union until the date of interview grouped into five-year groups, irrespective of whether the respondent is still married to her first partner.

Var Model Description

Sexual intercourse

Variables V525 to V528 relate to age at first intercourse, frequency of intercourse and time since last sexual relations. BASE (for variables ~~V526~~V527 to ~~V533~~V532): Respondents who have had sexual intercourse (V525 <> 0).

- V525 Age at first sexual intercourse. Respondents who had never had sex are coded 0. The response category "First sexual intercourse at first union" has been added in DHS III.
- V527 Time since the last sexual relations as reported by the respondent. The first digit gives the units in which the respondent gave her answer: 1 - Days ago, 2 - Weeks ago, 3 - Months ago, 4 - Years ago,, with 9 meaning a special answer was given. The last two digits give the time in the units given. Any value for time greater than 90 is a special answer.
- V528 Time since the last sexual intercourse in days is calculated from the preceding variable. Durations of more than 30 days are grouped into one category 31+. If the respondent said she had had sexual relations in the last four weeks, but replied that her last sexual intercourse was one month before the interview, then this is recoded to 30 days. Otherwise, one month is coded 31+ days.
- V529 Computed time since last sexual intercourse. This is computed from the responses for V527, with durations exceeding the interval since the last birth (V530 = 9) recoded as "Before last birth" and inconsistent responses flagged on variable V530 (codes 1-8) recoded to 97.
- V530 Flag variable indicating inconsistencies found in editing the responses for variable V527.
- 0 No flag
 - 1 Duration given is greater than the interval since the last birth and the respondent did not say, in the maternity section, that she was still abstaining from sexual relations since her last birth
 - 2 Duration given plus the duration of abstinence after the last birth is greater than the interval since the last birth
 - 3 Duration was reported, but the respondent had not resumed sexual intercourse since the last birth
 - 4 Respondent reported her last intercourse was before her last birth, but she had never given birth
 - 5 Respondent reported her last intercourse was before her last birth, but she was currently pregnant
 - 6 Respondent reported her last intercourse was before her last birth, but reported in the maternity section that she had resumed sexual intercourse after her last birth
 - 7 Respondent reported a duration since her last intercourse, but this would place her last intercourse before her last pregnancy
 - 8 Respondent reported a duration since her last intercourse, but this response was inconsistent with her response concerning the number of times she had had sexual intercourse in the four weeks preceding the survey.
 - 9 Duration was reported, but the duration would place her last sexual intercourse before her last birth.
- V531 Age at first sexual intercourse - imputed. This is the same as V525, except for respondents who reported that their first sexual intercourse was at the time of their union. For these cases, the age at first sex is taken from the age at first union. In cases where the age at first sex was inconsistent with the age at conception of the first child, but only by one year (V532

Var Model Description

= 3), the age at first sex was reduced by one year, consistent with the "Rule of one" applied in DHS I. Other cases flagged as inconsistent on variable V532 (codes 1, 2, 4, 5) are recoded as 97 (inconsistent). Cases coded 6 on V532 are not changed.

V532 Flag variable for inconsistencies found in editing the responses for V525.

0	No flag
1	Respondent reported age at first sexual intercourse that exceeds her current age
2	Respondent reported her age at first sexual intercourse as occurring more than one year <u>after</u> the conception of her first child
3	Respondent reported her age at first sexual intercourse as occurring up to one year <u>after</u> the conception of her first child
4	Respondent reported that her first sexual intercourse was at the time of her first marriage, but the respondent was never married
5	Respondent reported that her first sexual intercourse was at the time of her first marriage, but her first marriage occurred <u>after</u> the conception of her first child
6	Respondent reported her first sexual intercourse as being <u>after</u> her first marriage

~~V534 Whether the respondent, who is not currently married or living with a man, has a regular, occasional, or no sexual partner.
BASE: Women who are not currently married and not living with a man (V502 <= 1).~~

V535 Whether the respondent has ever been married or lived with a man.
BASE: Women who are not currently married and not living with a man (V502 <= 1).

V536 Recent sexual activity. It gives the sexual activity of the respondents during the last four weeks coded as follows:

- | | |
|---|--|
| 0 | Never had intercourse |
| 1 | Active in last 4 weeks |
| 2 | Not active in last 4 weeks - postpartum abstinence after the birth of the last child |
| 3 | Not active in last 4 weeks - not postpartum abstinence after the birth of the last child |

BASE: All women. Respondent who never had sex were coded 0.

V537 Months of abstinence. Postpartum or not postpartum abstinence.
BASE: Women sexually active during the last 4 weeks (V536 = 2 or V536 = 3).

Var Model Description

Section 61 (REC61)

Fertility Preferences

Var Model Description

V602 Fertility preferences. This variable comes primarily from a single question in the DHS III and MEASURE *DHS+* questionnaires. This is the same question used in the DHS I Model "B" questionnaire, and the DHS II and DHSIII Model "A" and "B" questionnaires. However, for DHS I Model "A" questionnaires, this variable was constructed from a series of questions. Women who respond that they want another child, but when asked when they would like the next child, respond that they cannot get pregnant, are classified in the "declared infecund category", and not in the "Wants another" category. These women can be identified in variable V616, where the original response to the question asking how long they would like to wait before having another child is recorded. In some countries, women who had never had sexual intercourse were not asked the questions relating to desire for future children, and are coded 6 on V602.

BASE: *All women.*

V603 Preferred waiting time before the birth of another child is created from a single question asking how long from the date of interview the respondent would like to wait before the birth of the next child. If the respondent answered "Don't know" or gave an "Other" answer when she was asked how long she would like to wait for her next child, she is no longer asked how old she would like her youngest child to be when the next child is born. In some countries there may be some additional non-numeric responses to the question of how long to wait before the next birth. These are assigned additional codes on a country-specific basis.

BASE: *All women who want another child (V602 = 1).*

V604 The preferred waiting time to the next birth is grouped into 12-month categories with responses of more than six years coded as 6+ years. Non-numeric responses are coded into one group (7 "Non-numeric"), but with "Don't know" and missing responses in their own categories (8 & 9). The additional response "Soon/Now" is not grouped with the other non-numeric codes, but is recoded as less than one year waiting time.

BASE: *All women who want another child (V602 = 1).*

V605 Desire for more children is a constructed variable classifying respondents who want more children by whether they want the next child soon (less than 2 years) or they want the next child later (2+ years). Sterilized women and women who want no more children are now recorded in separate categories. In some countries, women who had never had sexual intercourse were not asked the questions relating to desire for future children, and are coded 8 on V605.

BASE: *All women.*

~~V606 Respondent's attitude towards becoming pregnant.
BASE: *Non-pregnant, non-sterilized women (V213 = 0 & V312 < 6 & V312 < 7).*~~

V610 Whether the respondent thinks her partner approves of couples using a method to avoid pregnancy.

BASE: *Currently married or in union, non-sterilized women (V502 = 1).*

V611 How often the respondent discussed family planning with her partner in the past year.

<u>Var</u>	<u>Model</u>	<u>Description</u>
		BASE: Currently married or in union women (V502 = 1).
V612		Whether the respondent approves, in general, of couples using a method to avoid pregnancy.
V613		The ideal number of children that the respondent would have liked to have in her whole life, irrespective of the number she already has. In many countries it was possible for a respondent to reply to this question with a range of values, in which case this variable contains the midpoint between these values. If the midpoint is not an exact number then the number is rounded up in half the cases and rounded down for the other half. In situations where a range of values was collected, the original variables are included as country-specific variables. In some countries, additional country-specific categories are included, such as "It depends on God" or "As many as I can support" and are given country-specific codes.
V614		This variable groups the preceding variable such that 6 or more children are in one category 6+ and all non-numeric responses are coded 7.
V616		This variable records the original response to the question "How long would you like to wait from now before the birth of another child?" The first digit gives the units in which the respondent answered (1 indicates months, 2 indicates years, and 9 indicates a special response), while the last two digits give the time in those units. If the units value is 9 then the variable contains a special response, and if the duration value is greater than 90 this also indicates a special response. For example, code 994 is used for the responses "Soon/Now", and code 299 would mean that the response was given in years but the actual duration was missing on the questionnaire. BASE: All women who want another child (V602 = 1), plus those originally responding that they want another child, but then say they cannot get pregnant (see also V602).
V618		Regret for sterilization records whether the respondent regretted the sterilization and, if so, the reason she regretted the sterilization. The category child died has been added to this variable. BASE: Women who are sterilized or whose partners are sterilized (V312 = 6 or V312 = 7).
V621		Whether the respondent believes her partner wants the <u>same</u> number of children, <u>more</u> children or <u>fewer</u> children than she wants herself. BASE: Currently married or in union- women and neither partner is sterilized (V502 = 1 & V312 <> 6 & V312 <> 7).

Var Model Description

V623 The exposure status variable differentiates between pregnant women, postpartum amenorrheic women, menopausal or infecund women, and fecund women:

- Pregnant women.
- Postpartum amenorrheic women are those whose period has not returned since the last birth in the three/five years preceding the survey.
- Women are defined as being menopausal if they are not pregnant and not postpartum amenorrheic, are not currently using a contraceptive method, and have not had a period in the six months preceding the survey or report that they are in menopause.
- Women are defined as being infecund if they are not menopausal and not postpartum amenorrheic and not pregnant, have had no birth in the five years preceding the survey, and either (Model "A" countries) have been continuously married and have not used contraception in the five years preceding the survey, or (Model "B" countries) have been married one time and first married five or more years before the survey and have never used contraception.
- Fecund women are all women not included in the preceding categories.

V624 The need for family planning variable categorizes women according to whether they have an unmet need or a met need, to space or to limit their future births:

- Unmet need for spacing includes pregnant women whose pregnancy was mistimed, postpartum amenorrheic women whose last birth was mistimed, and fecund women who are neither pregnant nor postpartum amenorrheic and who are not using any method of family planning and say they want to wait two or more years for their next birth, are undecided about the timing of the next birth, or are undecided whether to have another child.
- Unmet need for limiting includes pregnant women whose pregnancy was unwanted, postpartum amenorrheic women whose last birth was unwanted and fecund women who are neither pregnant nor postpartum amenorrheic and who are not using any method of family planning and who want no more children.
- Met need for spacing includes women who are using some method of family planning and say they want to have another child, are undecided about the timing of the next birth, or are undecided whether to have another child.
- Met need for limiting includes women who are using family planning and who want no more children. Note that the specific methods are not taken into account here.

In Model "A" countries, pregnant and postpartum amenorrheic women whose pregnancy was the result of a contraceptive failure are not included in the category of unmet need, but are categorized as spacing failures or limiting failures. In Model "B" countries, no distinction is made since the information on contraceptive failure is not ascertained.

For formerly married and never married women, two additional categories exist. Women who have never had sex are separated into a separate category, as are women who would be categorized as having an unmet need to space or to limit, but who had not had sex in the month before the interview.

The remaining cases are those women who have no need for contraceptive methods, either because they desire a child soon (within the next two years) or because they are menopausal or infecund. Note that the infecund or menopausal category on this variable contains fewer cases than variable V623 as those women that are categorized as infecund or menopausal, but are currently using a contraceptive method are recorded in the two "met need" categories. Additionally, the code for the category "infecund or menopausal" has been changed to code 9 to allow for the two extra coding categories for formerly married or never married women.

<u>Var</u>	<u>Model</u>	<u>Description</u>
		BASE: All women. NOTE: This definition was used in the majority of the DHS II survey reports.
V625		Exposure status (definition 2) reclassifies variable V623, using a more liberal definition of infecundity. There are two differences between this definition and the definition used in V623: 1) For Model "B" countries, it is only possible to say that a women had been continuously married throughout the preceding five years if she was in her first union. This definition has been relaxed in V625, such that the respondent need only have been first married at least five years ago, and not necessarily continuously married throughout the last five years. For Model "A" countries, there is no change to this part of the definition. 2) Two additional variables have been used to declare a woman infecund. If the respondent said she cannot get pregnant when asked about preferences for additional children (V602 = 5), or if she reported that she was menopausal or had a hysterectomy when giving the reason she was not currently using a contraceptive method (V376 = 14), the respondent is coded as infecund.
V626		Unmet need (definition 2) follows exactly the same logic as V624, but uses the definition of fecundity given in V625. This variable was not used in DHS II survey reports, but has been used in the majority of DHSIII survey reports. BASE: <i>All women.</i>
V627		Ideal number of boys.
V628		Ideal number of girls.
V629		Ideal number of either sex. These three variables should sum to the total ideal number of children given in variables V613. If the response to the question for variables V613 is a non-numeric response, these variables are coded with the same response. In addition, there may be non-numeric responses on each of these questions. Country specific categories for non-numeric responses may also be recorded for these variables.
		Whether the respondent discussed the practice of family planning with any of the following people:
V630A		Husband or partner
V630B		Mother
V630C		Father
V630D		Sister(s)
V630E		Brother(s)
V630F		Daughter(s)
V630G		Mother-in-law
V630H		Friends or neighbors
V630I		<i>Country specific</i>
V630J		<i>Country specific</i>
V630K		<i>Country specific</i>
V630L		<i>Country specific</i>
V630M		Country specific
V630N		Country specific
V630O		Country specific
V630X		Other people
V631		If the respondent discovered that she became pregnant would that be a problem.

Var Model Description

V632 Women using contraception are asked who decided on the use of contraception.

Reason for not having sex

All respondents were asked about when they think is justified to refuse sex with their husband.

V633A Husband has STD
V633B Husband has other women
V633C Recent birth
V633D Tired, mood
V633E Country specific
V633F Country specific
V633G Country specific

Section 71 (REC71)

Partner's Characteristics and Women's Work

<u>Var</u>	<u>Model</u>	<u>Description</u>
V701		The current or most recent husband or partner's highest level of education attended. See variable V106. BASE: Ever-married women (V501 \diamond 0).
V702		Highest year of education gives the years of education completed at the level given in V701. BASE: Ever-married women except those answering "No education" or with missing data for V701 (V501 \diamond 0 & V701 \diamond 0 & V701 \diamond 8 & V701 \diamond 9).
V704		Current or last husband or partner's most recent occupation as collected in the country. Codes are country-specific. BASE: Ever-married women (V501 \diamond 0).
V705		Standardized partner's occupation groups. Agricultural categories also include fishermen, foresters and hunters and are <u>not</u> the basis for selection of agricultural/non-agricultural workers for the variables that follow. This selection is based on a country specific coding scheme in variable V704. In countries, where it is not possible to differentiate between self-employed agricultural workers and agricultural employees, no attempt has been made to use other information, and code 4 has been used for both categories. The analyst may wish to use other related information to differentiate between these two categories. BASE: Ever-married women (V501 \diamond 0).
V707		Whether the husband/partner works on his own land, /family land, rented land or on someone else's land. In DHS III, his own land and family land are differentiated between. BASE: Ever-married women whose partner works or worked in an agricultural occupation (V501 <math>\diamond</math> 0 & V704 = country-specific agricultural category).
V714		Whether the respondent is currently working.
V715		Most recent husband or partner's education in single years. See variable V133. BASE: Ever-married women (V501 \diamond 0).
V716		Respondent's occupation as collected in the country. Codes are country-specific. BASE: Women who are currently working or who have worked in the last 12 months (V731 = 1 or V731 = 2).
V717		Standardized respondent's occupation groups. Agricultural categories also include fishermen, foresters and hunters and are <u>not</u> the basis for selection of agricultural/non-agricultural workers. In countries, where it is not possible to differentiate between self-employed agricultural workers and agricultural employees, no attempt has been made to use other information, and code 4 has been used for both categories. The analyst may wish to use other related information to differentiate between these two categories. BASE: Women who are currently working or who have worked in the last 12 months (V731 = 1 or V731 = 2).
V718		Current type of employment. This variable is constructed from responses to the questions concerning who the respondent works for, whether she earns cash for this work, and whether she works at home or away from home.

<u>Var</u>	<u>Model</u>	<u>Description</u>
V719		Whether the respondent works for a family member, for someone else or is self-employed. BASE: Women currently working (V731 = 1 or V731 = 2).
V720		Whether the respondent received cash for this work. BASE: Women currently working (V731 = 1 or V731 = 2).
V721		Whether the respondent works at home or away from home. BASE: Women currently working (V731 = 1 or V731 = 2).
V723		Who usually takes care of the youngest child while the respondent is working. Women who have the child with them at work are coded 0. Coding categories for DHS III are slightly different from those used in DHS II. BASE: Women currently working or who have worked in the last 12 months and who have at least one living child aged 5 or less and still living at home (V731 = 1 or V731 = 2) & count(REC21 where B8 <= 5 & B5 = 1 & B9 = 0) > 0).
V729		Educational achievement recodes the education of the partner into the following categories: None, incomplete primary, complete primary, incomplete secondary, complete secondary, higher education, unknown level of education. If the grade within a level is unknown, it is assumed that the level was not completed. See related variables V702, V703, V715. BASE: Ever-married women (V501 > 0).
V730		Age of the respondent's husband or partner. BASE: Currently married or in union women (V502 = 1).
V731		Whether the respondent worked in the last 12 months.
V732		Whether the respondent works throughout they year, seasonally, or just occasionally. BASE: Women who are currently working or who have worked in the past year (V731 = 1 or V731 = 2).
V733		For seasonal or part year workers, the number of months they worked in the last twelve months. BASE: Women who are working seasonally or for part of the year (V732 = 2).
V734		Number of days the respondent usually worked per week. BASE: Women who either work throughout the year or seasonally (V732 = 1 or V732 = 2).
V735		The approximate number of days the respondent worked in the last twelve months. For respondents who worked throughout the year, this is 50 times the usual number of days worked per week. For respondents who worked for part of the year or who worked seasonally, this is the number of months worked times the number of days usually worked per week times 50/12. For women who only work occasionally, this is the number of days worked in the last twelve months, as reported in the questionnaire. BASE: Women who are currently working or who worked in the past year (V731 = 1 or V731 = 2).
V736		Usual amount the respondent earns in cash for the work she does. This variable is 8 digits in size. The first digit gives the units in which the amount was specified, while the remaining digits give the total amount. The first digit, or units digit, is coded as follows: — 1 — per hour — 2 — per day

<u>Var</u>	<u>Model</u>	<u>Description</u>
		<p>— 3 — per week</p> <p>— 4 — per month</p> <p>— 5 — per year</p> <p>For example, 30000400 would indicate that the respondent received 400 per week. The currency and units of currency used are country specific.</p> <p>BASE: Women paid cash for their work (V720 = 1).</p>
V737		Approximate daily earnings.
V738		Approximate yearly earnings.
		<p>These two variables are calculated from the preceding variables, in an attempt to standardize earnings to common units. The currency and currency units used are country specific.</p> <p>BASE: Women paid cash for their work (V720 = 1).</p>
V739		<p>The person who mainly decides how the money earned by the respondent is used.</p> <p>BASE: Women paid cash for their work (V720 = 1).</p>
V740		<p>Whether the respondent works on her own land, family land, rented land or on someone else's land.</p> <p>BASE: Women who are currently working or who have worked in the last 12 months, and who work or worked in agriculture (V716 = country-specific agricultural category).</p>
V741		<p>Type of earnings for work. The respondent is asked whether they receive cash for their work, they are paid in kind, a combination of the two or not paid.</p> <p>BASE: Respondents who have worked during the last 12 months (V731 = 1 or V731 = 2).</p>
V742		<p>Portion of household expenditures respondents earnings pay.</p> <p>BASE: Respondents earning cash for their work (V741 = 1 or V741 = 2).</p>
		<u>Final say in the family on the following decisions</u>
V743A		Respondent's health care
V743B		Making large household purchases
V743C		Making household purchases for daily needs
V743D		Visits to family or relatives
V743E		Food to be cooked each day
		<u>When wife's beating or hitting is justified</u>
V744A		Goes out without telling him
V744B		Neglects the children
V744C		Argues with him
V744D		Refuses to have sex with him
V744E		Burns the food.

Section 75 (REC75)

AIDS and Condom Use

<u>Var</u>	<u>Model</u>	<u>Description</u>
V750		Heard of AIDS or other sexually transmitted Diseases.
V751		Whether the respondent has ever heard of AIDS (Acquired Immune Deficiency Syndrome).
		Sources of information from which the respondent has learned most about AIDS.
V752A	Radio	Radio
V752B	Television	Television
V752C	Newspapers or magazines	Newspapers or magazines
V752D	Pamphlets or posters	Pamphlets or posters
V752E	Clinics or health workers	Clinics or health workers
V752F	Churches or mosques	Churches or mosques
V752G	Schools or teachers	Schools or teachers
V752H	Community meetings	Community meetings
V752I	Friends or relatives	Friends or relatives
V752J	Work place	Work place
V752K	Country specific	Country specific
V752L	Country specific	Country specific
V752M	Country specific	Country specific
V752N	Country specific	Country specific
V752O	Country specific	Country specific
V752X	Other responses	Other responses
		BASE: Women who have heard of AIDS (V751 = 1).
V753		Whether the respondent believes there is anything a person can do to avoid AIDS. BASE: Women who have heard of AIDS (V751 = 1).
		Ways in which the respondent thinks people can avoid AIDS
V754A		"Safe Sex"
V754B		Abstaining from sex
V754C		Using condoms during sex
V754D		Having only one sexual partner
V754E		Avoiding sex with prostitutes
V754F		Avoiding sex with homosexuals
V754G		Avoiding blood transfusions
V754H		Avoiding injections
V754I		Avoiding kissing
V754J		Avoiding mosquito bites
V754K		Seeking protection from a traditional healer
V754L		<i>Limit number of sexual partners</i>
V754M		<i>Avoid partners who have many partners</i>
V754N		<i>Avoid sex with intravenous drug users</i>
V754O		<i>Avoid sharing razor blades with AIDS patients</i>
V754P		<i>Country specific</i>
V754Q		<i>Country specific</i>
V754R		<i>Country specific</i>
V754S		<i>Country specific</i>
V754T		<i>Country specific</i>

V754U *Country specific*
 V754V *Country specific*
 V754X Other responses
 V754Z Does not know any means of avoiding AIDS
 BASE: Women who have heard of AIDS (V751 = 1).

Reduce chances of AIDS

Respondents were asked whether using condoms or having just one sexual partner would reduce their chances of getting AIDS.
 BASE: Heard of AIDS and there are ways to avoid AIDS (V751 = 1 | V753 = 1)

V754CP Always using condoms during sex
 V754DP 1 sex partner with no other partners

Get AIDS

Respondents were asked if transmission of AIDS can happen by being bitten by a mosquito or by sharing food with a person who has AIDS.
 BASE: Heard of AIDS and there are ways to avoid AIDS (V751 = 1 | V753 = 1)

V754JP From mosquito bites
 V754WP Sharing food with person who has AIDS

What "Safe Sex" means to the respondent

~~V755B Abstaining from sex~~
~~V755C Using condoms during sex~~
~~V755D Having only one sex partner~~
~~V755E Avoiding sex with prostitutes~~
~~V755F Avoiding sex with homosexuals~~
 V755L *Country specific*
 V755M *Country specific*
 V755N *Country specific*
 V755O *Country specific*
 V755X Other responses
 V755Z Does not know the meaning of "Safe sex"
 BASE: ~~Women who responded that "Safe sex" was a way of avoiding AIDS (V754A = 1).~~

V756 Whether the respondent believes it is possible for a healthy-looking person to have the AIDS virus.
 BASE: Women who have heard of AIDS (V751 = 1).

~~V757 Whether the respondent believes that AIDS is a fatal disease.~~
 BASE: ~~Women who have heard of AIDS (V751 = 1).~~

~~V758 Whether the respondent believes her risk of getting AIDS is small, moderate, great, no risk at all, or that she already has AIDS.~~
 BASE: ~~Women who have heard of AIDS (V751 = 1).~~

Whether it is acceptable to discuss AIDS on the following media:

BASE: Heard of AIDS (V751 = 1)

V759A X Radio

Var Model Description

V759B X Television
 V759C X Newspapers
 V759D X Country Specific
 V759E X Country Specific
 V759F X Country Specific
 V759G X Country Specific
 V759H X Country Specific
 V759I X Country Specific

~~Ways in which the respondent has changed her sexual behavior, since hearing about AIDS, in order to avoid getting AIDS:~~

~~V760A Did not start sex
 V760B Stopped all sex
 V760C Started using condoms during sex
 V760D Restricted the number of partners to one
 V760E Reduced the number of partners
 V760F Ask spouse to be faithful
 V760G No more homosexual contacts
 V760I Stopped receiving injections
 V760L Country specific
 V760M Country specific
 V760N Country specific
 V760O Country specific
 V760P Ask spouse to avoid prostitutes
 V760V No non-sexual change in behavior
 V760W Other (non-sexual) responses
 V760X Other (sexual) responses
 V760Y Did not change behavior
 V760Z Don't know whether they changed behavior~~

~~BASE: Women who have heard of AIDS and have ever had sexual intercourse (V751 = 1 & V525 <> 0).~~

V761 Whether the respondent used a condom the last time she had sexual intercourse.
 BASE: Women who have ever had sexual intercourse (V525 <> 0).

Used condom during intercourse

V761B With other man (1)
 V761C X With other man (2).

~~V762 Source of condoms known by the respondent. If the respondent does not know where to get condoms, this variables is coded 98.~~

~~BASE: Women who have ever had sexual intercourse (V525 <> 0).~~

Knowledge of a source for male condoms

V762AA Government hospital
 V762AB Government health center/post
 V762AC Family planning clinic
 V762AD Mobile clinic
 V762AE Public field worker

<u>Var</u>	<u>Model</u>	<u>Description</u>
V762AF		Other public
V762AG		Public Country specific
V762AH		Public Country specific
V762AI		Public Country specific
V762AJ		Private hospital, clinic
V762AK		Pharmacy
V762AL		Private doctor
V762AM		Private mobile clinic
V762AN		Private field worker
V762AO		Other private
V762AP		Private Country specific
V762AQ		Private Country specific
V762AR		Private Country specific
V762AS		Shop
V762AT		Church
V762AU		Friends, relatives
V762AV		Country specific
V762AW		Country specific
V762AX		Other
V762AZ		Don't know a source for condom

Knowledge of a source for female condoms

V762BA		Government hospital
V762BB		Government health center/post
V762BC		Family planning clinic
V762BD		Mobile clinic
V762BE		Public field worker
V762BF		Other public
V762BG		Public Country specific
V762BH		Public Country specific
V762BI		Public Country specific
V762BJ		Private hospital, clinic
V762BK		Pharmacy
V762BL		Private doctor
V762BM		Private mobile clinic
V762BN		Private field worker
V762BO		Other private
V762BP		Private Country specific
V762BQ		Private Country specific
V762BR		Private Country specific
V762BS		Shop
V762BT		Church
V762BU		Friends, relatives
V762BV		Country specific
V762BW		Country specific
V762BX		Other
V762BZ		Don't know a source for condom

Sexually transmitted diseases (STD) caught in last 12 months

Var Model Description

V763A X Any STD
 V763B X Genital sore/ulcer
 V763C X Genital discharge
 V763D X Country specific
 V763E X Country specific
 V763F X Country specific
 V763G X Country specific

~~V764 Whether the respondent has ever heard of condoms for contraceptive use or for use to prevent STDs.~~

~~V765 Whether the respondent has ever used condoms for contraceptive use or for use to prevent STDs.~~

~~Both of these variables are created from responses to several questions in the questionnaire, but the set of questions used may vary from country to country, depending on the depth of questioning used in a particular country.~~

Reason of condom use during last sexual intercourse

V765A X Last time
 V765B X With another man (1)
 V765C X With another man (2)

Number of sexual partners during the last 12 months

V766A Number of men other than the husband
 V766B Number of men including the husband

Relationship with last sexual partner

V767A Last partner
 V767B Other sexual partner (1)
 V767C Other sexual partner (2)

Length of time knows last sexual partner

V768A Last partner
 V768B Other sexual partner (1)
 V768C Other sexual partner (2)

Could get a condom herself

V769 Male condom
 V769A Female condom

Seek advice or treatment for an STD

V770 X Last STD infection

Place where advice or treatment is sought

Var Model Description

V770A X Clinic/hospital/private doctor
 V770B X Traditional healer
 V770C X Shop/pharmacy
 V770D X Friends/relatives
 V770E X Country specific
 V770F X Country specific
 V770G X Country specific
 V770H X Country specific
 V770I X Country specific

V771 X Advise partner when had disease
 V772 X Tried to avoid infecting partner

What done to prevent infecting the sexual partner

V773A X Abstaining from sexual intercourses
 V773B X Used condoms
 V773C X Took medicines
 V773D X Country specific
 V773E X Country specific
 V773F X Country specific
 V773X X Other

Mother to child AIDS transmission

V774 AIDS can be transmitted from mother to child

Ways of transmission from mother to child

V774A During pregnancy
 V774B During delivery
 V774C By breastfeeding

V775 Respondent knows personally someone who has or died of AIDS

V776 Respondent spoke with spouse about avoiding AIDS

V777 Family allowed to keep secret an AIDS infection of a member

V778 Respondent willing to care for a relative in the household who has AIDS

V779 X Person with AIDS allowed to continue teaching

V780 X Should children be taught about condoms

AIDS test

V781 X Ever been tested for AIDS
 V782 X Want to be tested for AIDS
 V783 X Know a place to get AIDS test

Var Model Description

V784 X Place where someone can get an AIDS test

V785 Heard about other STDs

Man's symptoms of sexually transmitted infections (STIs)

BASE: Women who have heard about other STDs (V785 = 1).

V785A Abdominal Pain
 V785B Genital discharge or dripping
 V785C Foul smelling discharge
 V785D Burning pain on urination
 V785E Redness, inflammation of the genital area
 V785F Swelling in the genital area
 V785G Genital sores or ulcers
 V785H Genital warts
 V785I Genital itching
 V785J Blood in urine
 V785K Loss of weight
 V785L Impotence
 V785M Country specific
 V785N Country specific
 V785O Country specific
 V785P Country specific
 V785Q Country specific
 V785X Other
 V785Y No symptoms
 V785Z Don't know

Woman's symptoms of sexually transmitted infections (STIs)

BASE: Women who have heard about other STDs (V785 = 1).

V786A Abdominal pain
 V786B Genital discharge
 V786C Foul smelling discharge
 V786D Burning pain on urination
 V786E Redness or inflammation of the genital area
 V786F Swelling in the genital area
 V786G Genital sores or ulcers
 V786H Genital warts
 V786I Genital itching
 V786J Blood in urine
 V786K Loss of weight
 V786L Hard to get pregnant or to have a child
 V786M Country specific
 V786N Country specific
 V786O Country specific
 V786P Country specific
 V786Q Country specific
 V786X Other
 V786Y No symptoms
 V786Z Don't know

Var Model Description

Alcohol drinking during the last 3 months

V787 X Number of days drank alcohol

V788 X Number of days gotten drunk

Injections received last 3 months

V789 X Number of injections in last 3 months

V790 X Person who gave last injection

Section 81 (REC81)

Characteristics of the Interview

<u>Var</u>	<u>Model</u>	<u>Description</u>
V801		Time of the start of the interview. The first two digits give the time in hours using the 24-hour clock, and the last two digits give the minutes within that hour.
V802		Time of the end of interview is coded as for the start of interview.
V803		Length of interview in minutes is calculated from the previous two variables, but with interviews that required more than one visit being coded 96.
V804		Number of visits for the interview.
V805		Interviewer identification code. Codes are country-specific. This variable occupies 3 digits starting from DHS III.
V806		Data entry keyer code. Codes are country-specific.
V811		Presence of children aged under 10 at the end of the husband's background and woman's work section of the interview.
V812		Presence of the husband at the end of the marriage and sexual intercourse section of the interview.
V813		Presence of other males at the end of the marriage and sexual intercourse section of the interview.
V814		Presence of other females at the end of the marriage and sexual intercourse section of the interview.

Section 82 (REC82)

Calendar

Var Model Description

VCOL A Column number of the entries in the calendar, indicating the type of data found in the entry in the calendar.

VCAL A The calendar of events representing the 5+ years prior to the date of interview. The calendar is split into 5 records, representing each of the 5 columns. Each of the 5 columns contains a single character for each month in the time period. The data are stored as single variables of 80 characters, allowing for up to 80 months to be represented in the calendar. The first character in each variable represents the most recent point in time, while the 80th character position represents data for January of the year in which the calendar started. The calendars are fixed at the 80th character position, such that the first few entries in the calendar represent points in time after the date of interview, and are consequently left blank. The columns are as follows:

Column Description

- | | |
|---|--|
| 1 | Births, pregnancies and contraceptive use |
| 2 | Reasons for discontinuation of contraceptive use |
| 3 | Marital/union status |
| 4 | Country specific |
| 5 | Source of contraception |

The entire calendar is usually inserted in questionnaires of countries using the model A. However, MEASURE *DHS+* includes, in model B questionnaires, a reduced calendar that has one column on births, pregnancies and contraceptive use (column 1).

Column Description

- 1 Records each of the births and pregnancies during the calendar period, as well as each episode of contraceptive use and non-use. The following codes are used in column 1:
- | | | | |
|---|----------------------------|----------|-------------------------------------|
| 0 | Non-use of contraception | L | Lactational amenorrhea |
| 1 | Pill | C | Female condom |
| 2 | IUD | F | Foam and Jelly |
| 3 | Injections | α | Country-specific method 1 |
| 4 | Diaphragm | β | Country-specific method 2 |
| 5 | Condom | τ | Country-specific method 3 |
| 6 | Female sterilization | ? | Unknown method/missing data |
| 7 | Male sterilization | B | Birth |
| 8 | Periodic abstinence/rhythm | T | Terminated pregnancy/non-live birth |
| 9 | Withdrawal | P | Pregnancy |
| W | Other traditional methods | | |
| N | Norplant | | |
| A | Abstinence | | |

All codes are standard except for codes α , β , and τ which are country-specific letter codes representing traditional methods.

- 2 Records the reason for discontinuation of a method. The discontinuation code appears in the row of the last month of use of the method for the episode. All other rows in the column are

Column

Description

left blank, except for those in which discontinuations took place. The standard codes are as follows:

1	Became pregnant while using	C	Cost
2	Wanted to become pregnant	F	Fatalistic
3	Husband disapproved	A	Difficult to get pregnant/menopause
4	Side effects	D	Marital dissolution
5	Health concerns	W	Other reasons
6	Access/availability	K	Don't know
7	Wanted more effective method	α	Country-specific reason 1
8	Inconvenient to use	β	Country-specific reason 2
9	Infrequent sex/husband away	τ	Country-specific reason 3

All codes are standard except for codes α , β , and τ which are country-specific letter codes representing additional reasons for discontinuation.

- 3 Records the episodes of marriage in the calendar period. The following codes are used:
- | | |
|---|---------------------------------------|
| X | In union (married or living together) |
| 0 | Not in union |
- 4 Country specific
- 5 Source of contraception

Rows in the calendar, representing months after the month of interview, are left blank. With this exception, columns 1, 3 and 4 do not contain any blank characters.

Var Model Description

Section 83 (REC83)

Maternal Mortality

The Maternal Mortality section is a country specific section that exists only for those countries that have a maternal mortality module. It contains up to 20 entries containing information related to all of the sisters and brothers of the respondents.

Var Model Description

MMIDX	MM	Index to maternal mortality history.
MM1	MM	Sex of sibling.
MM2	MM	Whether the sibling is dead or alive.
MM3	MM	Current age of sibling in years. BASE: All living siblings (MM2 = 1).
MM4	MM	CMC date of birth of sibling. This is based on a crude imputation process. The analysts may elect to perform their own imputation based on their own assumptions.
MM5	X	Sibling's marital status. Whether the sibling is/was ever married or not. BASE: All siblings whose age is/was greater than or equal to the cut off age (MM3 >= MMC5).
MM6	MM	Number of years ago the respondent's brother or sister died.
MM7	MM	Age at death of sibling in years.
MM8	MM	CMC date of death of sibling. The analyst may choose to perform their own imputation procedure, as for the CMC date of birth of the sibling. BASE: All siblings who have died (MM2 = 0).
MM9	MM	Indicates if the respondent's sister was pregnant when she died, if she died during childbirth, within six weeks after the delivery, within 2 months after the delivery. BASE: Female siblings aged older than the cutoff age (MM1 = 2 & MM2 = 0 & MM7 >= MMC5). In some countries the question is only asked for ever married siblings (MM5 = 1).
MM10	MM	Information about whether the death that occurred was related to the sister's pregnancy. BASE: Female siblings that died during a pregnancy or a period of time after a delivery or a still birth (MM9 = 2 or MM9 = 4 or MM9 = 5 or MM9 = 6).
MM11	X	Specifies the cause of death. Codes are country specific. BASE: All deaths not related to a pregnancy for a specified age and years within which death occurred (see original questionnaire for each country for further details).
MM12	X	Time between delivery and death -- used in countries where questions relating to fixed periods of time between delivery and death (usually six weeks or two months) are not asked. BASE: Female siblings who died after pregnancy (MM9 = 4 or MM9 = 5 or MM9 = 6).
MM13	X	Place in which the death occurred. Country specific codes. BASE: Siblings who have died (MM2 = 0).

Var Model Description

MM14	MM	Number of children to whom the female sibling gave birth during her live. In most countries, for women with a maternity related death, this is the number of children born prior to the pregnancy, if the respondent was pregnant when she died or if she died during delivery, and includes the child, if the woman died after the birth of the child. BASE: As for MM9 above.
MM15	MM	Year of death of sibling. BASE: All siblings who have died (MM2 = 0).

Section 84 (REC84)**Maternal Mortality**

<u>Var</u>	<u>Model</u>	<u>Description</u>
MMC1	MM	Number of occurrences of the maternal mortality section. This variable gives the number of the respondent's brothers and sisters born to the same mother including the respondent.
MMC2	MM	Number of births to the respondent's mother preceding the respondent's birth. This variables gives the number of brothers and sisters born to the respondent's mother who are/were older than the respondent.
MMC3	X	In some countries information about the siblings is given by one of the sisters of the respondent, and not by the respondent herself, if both the sibling and the respondent were interviewed. In this variable, the line number of the sibling providing the information is recorded, if the information was not given by the respondent. If the respondent gave the information for the maternal mortality section, this variables is coded 0. The information for the respondent is copied from that reported by the sister, replacing the respondent's data by here sister's data in the maternal mortality section.
MMC4	X	Line numbers in the household schedule of the eligible sisters of the respondent.
MMC5	MM	Cut off age for this section. In most countries the age of 12 was taken as the cut off age, however in some countries the age of 10, 13 or 15 was used.

Var Model Description

Last Sexual Intercourse, Condom Use and Other Partners

Variables V850 to V852 relate to the time since last sexual intercourse and whether condoms were used at that time, for sexual relationships with the respondent's husband or partner and with other people, respectively. Variable V852 gives the number of other partners.

BASE (for V850 to V852): Currently married or in union women (V502 = 1)

- V850 Time since the respondent's last sexual intercourse with her husband or partner. The first digit gives the units in which the respondent gave her answer: 1—Days ago, 2—Weeks ago, 3—Months ago, 4—Years ago, with 9 meaning a special answer was given. The last two digits give the time in the units given. Any value for time greater than 90 is a special answer.
- V850A Whether a condom was used the last time the respondent had sexual intercourse with her husband or partner.
- V851 Time since the last sexual intercourse with someone other than the respondent's husband or partner. Respondents who have had no other partner in the prior twelve months are coded 995.
- V852 Whether a condom was used the last time the respondent had sexual intercourse with someone other than her husband or partner.
- V853 Number of partners other than the husband or partner with whom the respondent lives, with whom the respondent had sexual intercourse in the 12 months prior to the interview.

Knowledge of Sexually Transmitted Diseases (STDs)

Variables V855 to V856Z relate to knowledge of sexually transmitted diseases.

BASE (for V856A to V856Z): Heard of any sexually transmitted disease (V855 = 1).

- V855 Whether the respondent has ever heard of any sexually transmitted diseases.
- V856A Knowledge of Syphilis
- V856B Knowledge of Gonorrhea
- V856C Knowledge of AIDS
- V856D Knowledge of Genital warts
- V856E Knowledge of country specific
- V856F Knowledge of country specific
- V856G Knowledge of country specific
- V856H Knowledge of country specific
- V856I Knowledge of country specific
- V856J Knowledge of country specific
- V856K Knowledge of country specific
- V856X Knowledge of other diseases
- V856Z Does not know sexually transmitted diseases by name.

Var Model Description

Prevalence of STDs

Variables V857 to V858Z relate to whether the respondent has had any sexually transmitted disease in the twelve months prior to the interview.

BASE (for V858A to V858Z): Whether the respondent has ever had a sexually transmitted disease in the twelve months prior to the interview (V857 = 1).

V857 ————— Whether the respondent has had a sexually transmitted disease in the twelve months prior to the interview.

BASE: Women who had ever had sexual intercourse and had heard of sexually transmitted diseases (V525 < 0 & V855 = 1).

- V858A ————— Last 12 months had syphilis
- V858B ————— Last 12 months had gonorrhea
- V858C ————— Last 12 months had AIDS
- V858D ————— Last 12 months had genital warts
- V858E ————— Last 12 months had *country specific*
- V858F ————— Last 12 months had *country specific*
- V858G ————— Last 12 months had *country specific*
- V858H ————— Last 12 months had *country specific*
- V858I ————— Last 12 months had *country specific*
- V858J ————— Last 12 months had *country specific*
- V858X ————— Last 12 months had other disease
- V858Z ————— Last 12 months had unknown disease

Treatment of STDs

Variables V859 to V860Z relate to treatment of sexually transmitted diseases (STDs).

BASE (for V860A to V860Z): Women who sought advise for the last sexually transmitted disease (V859 = 1).

V859 ————— Sought advise for the last sexually transmitted disease.
BASE: Women who had a sexually transmitted disease in the 12 months prior to interview (V857 = 1).

- V860A ————— Sought advice from a government hospital
- V860B ————— Sought advice from a government health center
- V860C ————— Sought advice from a health post or dispensary
- V860D ————— Sought advice from a mobile clinic
- V860E ————— Sought advice from a community health worker
- V860F ————— Sought advice from a family planning clinic
- V860G ————— Sought advice from *country specific* public sector
- V860H ————— Sought advice from *country specific* public sector
- V860I ————— Sought advice from other public sector sources
- V860J ————— Sought advice from a private hospital or clinic
- V860K ————— Sought advice from a private pharmacy
- V860L ————— Sought advice from a private doctor
- V860M ————— Sought advice from a private mobile clinic
- V860N ————— Sought advice from a community health worker
- V860O ————— Sought advice from *country specific* medical private sector

Var Model Description

V860P	Sought advice from <i>country specific</i> medical private sector
V860Q	Sought advice from <i>country specific</i> medical private sector
V860R	Sought advice from other private sector sources
V860S	Sought advice from a shop
V860T	Sought advice from a traditional practitioner
V860U	Sought advice from relatives or friends
V860V	Sought advice from <i>country specific</i> other sector
V860W	Sought advice from <i>country specific</i> other sector
V860X	Sought advice from other sources
V860Z	Sought advice from an unknown source

Avoidance of infecting others with STD

Variables V861 to V863X provide information on what the respondent did when she realized that she had a sexually transmitted disease.

BASE (for V863A to V863X): Whether the respondent tried to avoid infecting her partner (V863 = 1).

V861	Whether the respondent advised her partner when she had the sexually transmitted disease. BASE: Women who had a sexually transmitted disease in the twelve months prior to the survey (V857 = 1).
V862	Whether the respondent tried to avoid infecting her partner. BASE: Women who had a sexually transmitted disease in the twelve months prior to the survey (V857 = 1).

Means of avoiding infecting her partner:

V863A	No sexual intercourse
V863B	Used condoms
V863C	Took medicines
V863D	<i>Country specific</i>
V863E	<i>Country specific</i>
V863F	<i>Country specific</i>
V863W	Other (non-sexual)
V863X	Other (sexual)

Knowledge of Means of Transmission of AIDS

Variables V864A to V864Z provide information on the ways in which the respondent believes a person can get AIDS. Variables V865 to V867 relate specifically to whether AIDS can be cured, whether it is transmitted from mother to child, and whether the respondent knows anyone who has AIDS or who has died of AIDS.

BASE: Respondents who have heard of AIDS (V751 = 1).

V864A	Get AIDS from sexual intercourse
V864B	Get AIDS from sex with multiple partners
V864C	Get AIDS from sex with prostitutes
V864D	Get AIDS from not using a condom
V864E	Get AIDS from homosexual contact
V864F	Get AIDS from blood transfusions
V864G	Get AIDS from injections
V864H	Get AIDS from kissing

Var Model Description

V864I ————— Get AIDS from mosquito bites
 V864J ————— Get AIDS from *country specific*
 V864K ————— Get AIDS from *country specific*
 V864L ————— Get AIDS from *country specific*
 V864M ————— Get AIDS from *country specific*
 V864N ————— Get AIDS from *country specific*
 V864O ————— Get AIDS from *country specific*
 V864P ————— Get AIDS from *country specific*
 V864X ————— Get AIDS from: other responses
 V864Z ————— Get AIDS from unknown sources

V865 ————— Whether the respondent believes AIDS can be cured.
 V866 ————— Whether the respondent believes that AIDS can be transmitted from mother to child.
 V867 ————— Whether the respondent knows someone who has AIDS or who has died of AIDS.

Reasons Respondent Assesses Risk of Getting AIDS to be Low

Variables V868B to V868Z give the reasons the respondent believes that she is at no risk or at a small risk of getting AIDS.

BASE: Respondents reporting they are at no risk or at only a small risk of getting AIDS (V758 = 0 or V758 = 1).

V868B ————— No/small risk: abstains from sex
 V868C ————— No/small risk: uses condoms
 V868D ————— No/small risk: has only one sexual partner
 V868E ————— No/small risk: has a limited number of partners
 V868F ————— No/small risk: spouse has no other partners
 V868G ————— No/small risk: has no homosexual contact
 V868H ————— No/small risk: has not received a blood transfusion
 V868I ————— No/small risk: has had no injections
 V868J ————— No/small risk: country specific
 V868K ————— No/small risk: country specific
 V868L ————— No/small risk: country specific
 V868P ————— No/small risk: spouse avoids prostitutes
 V868X ————— No/small risk: other reasons
 V868Z ————— No/small risk: reasons unknown

Reasons Respondent Assesses Risk of Getting AIDS to be High

Variables V869C to V869Z give the reasons the respondent believes that she is at a moderate or great risk of getting AIDS.

BASE: Respondents reporting they are at a moderate or great risk of getting AIDS (V758 = 2 or V758 = 3).

V869C ————— Great/moderate risk: not using condoms
 V869D ————— Great/moderate risk: has more than one sex partner
 V869E ————— Great/moderate risk: has many sex partners
 V869F ————— Great/moderate risk: spouse has other sexual partners
 V869G ————— Great/moderate risk: has homosexual contacts
 V869H ————— Great/moderate risk: has received blood transfusions
 V869I ————— Great/moderate risk: has had injections

Var Model Description

V869J ————— Great/moderate risk: country specific
V869K ————— Great/moderate risk: country specific
V869L ————— Great/moderate risk: country specific
V869P ————— Great/moderate risk: spouse frequents prostitutes
V869X ————— Great/moderate risk: other reasons
V869Z ————— Great/moderate risk: reasons unknown

Condom Use in Relation to AIDS

V870 ————— Whether the respondent has heard of using condoms to avoid AIDS.

V871 ————— Whether the respondent has ever used a condom to avoid AIDS.
BASE: Women who reported having heard of using condoms to avoid AIDS and who had ever had sexual intercourse (V870 = 1 & V525 \diamond 0).

Payments or Gifts for Sexual Intercourse

V872 ————— Whether the respondent ever received or gave money or gifts in return for sexual intercourse.
BASE: Women who had ever had sexual intercourse (V525 \diamond 0).

Sections 91-99 (REC91-REC99)

Country-Specific Variables

The following sections will appear in the recode data file as needed on a country-specific basis.

- REC91 All single occurrence country-specific variables relating to the respondent.
- REC92 Country-specific variables from the birth history (REC21). Variable IDX92 is always included as the first variable in this section and is equal to BIDX for each entry in the birth history.
- REC94 Country-specific variables from the maternity history (REC41). Variable IDX94 is always included as the first variable in this section and is equal to MIDX for each entry in the maternity history.
- REC95 Country-specific variables from the health history (REC43). Variable IDX95 is always included as the first variable in this section and is equal to HIDX for each entry in the health history.
- REC96 Country-specific variables from the height and weight table (REC44). Variable IDX96 is always included as the first variable in this section and is equal to HWIDX for each entry in the height and weight table.
- REC97-99 The last three country-specific sections are not assigned to any particular section of the questionnaire, but are used for additional modules not usually incorporated in the questionnaires. These include the respondent's work history, the diagnoses of deaths for dead children who were born in the three/five years preceding the interview, pregnancy history, or for husband's questionnaires.
- Note: As the child related sections REC41, REC43, REC44 are now completely parallel, i.e. the first entry in each section relates to the last child born, the second entry in each section relates to the last but one child born, etc., country specific variables for these sections, usually placed in REC94, REC95 or REC96, may all be placed in REC94 if the number of variables involved is small. This is to save space in the data file.

Section and Variable Description - Household

<u>Section</u>	<u>Code</u>	<u>Length</u>	<u>Class</u>	<u>Occurrences</u>		<u>Section label</u>
				<u>Min</u>	<u>Max</u>	
RECH0	H0	129	S	1	1	Household's Basic Data
RECH1	H1	57	M	0	90	Household Schedule
RECH2	H2	56	S	0	1	Household Characteristics
RECH3	H3	?	S	0	1	Country-specific Household Variables
RECH4	H4	?	M	0	90	Country-specific Household Schedule
RECH5	H5	90	M	0	15	Women Height/Weight/Hemoglobin
RECH6	H6	116	M	0	15	Children Height/Weight/Hemoglobin

? implies that the entry is country-specific

Section H0 (RECH0)

Household's Basic Data

<u>Var</u>	<u>Model</u>	<u>Description</u>
HHID		Case identification uniquely identifies each household. In most surveys this is constructed by concatenating the cluster or sample point number and the household number, but in some surveys this may be the questionnaire number taken from the front page of the questionnaire.
HV000		Alphabetic country code to identify the survey from which the data were collected. The code is based on an international standard code. This variable is 3 characters in length, with the third character indicating the format of the recode file used for this survey. For all surveys in MEASURE <i>DHS+</i> following this standard, this code will be 4. For example: DR4 is the Dominican Republic, MA4 is Morocco, ZM4 is Zambia, and ID4 is Indonesia.
HV001		Cluster number is the number identifying the sample point as used during the fieldwork. This variable may be a composite of several variables in the questionnaire. If so, the original variables are included in RECH3 as country-specific variables.
HV002		Household number is the number identifying the household within the cluster or sample point. In some cases, this variable may be the combination of dwelling number and household number within dwelling. In these cases, the original variables are included as country-specific variables.
HV003		Respondent's line number is the line number in the household schedule of the person responding to the questions asked in the household questionnaire. If nobody in the household was available for interview, this variable is coded 00.
HV004		Ultimate area unit is a number assigned to each sample point to identify the ultimate area units used in the collection of data. This variable is usually the same as the cluster number, but may be a sequentially numbered variable for samples with a more complicated structure.
HV005		Sample weight is an 8 digit variable with 6 implied decimal places. To use the sample weight divide it by 1000000 before applying the weighting factor. All sample weights are normalized such that the weighted number of cases is identical to the unweighted number of households when using the full dataset with no selection. This variable should be used to weight all tabulations produced using the data file. For self-weighting samples this variable is equal to 1000000.
HV006		Month of interview.
HV007		Year of interview. The size of this variable was changed from 2 digits in the previous surveys to 4 digits in MEASURE <i>DHS+</i> .
HV008		Century month code of date of interview (see note on century month codes).
HV009		Total number of household members indicates the number of entries to be found in RECH1.
HV010		Total number of eligible women indicates the number of women found eligible for the individual survey in the household schedule. The eligibility criteria are generally: female, aged between 15 and 49. In some countries, the eligibility criteria restricts the survey to ever-married women. In early DHS II surveys, the eligibility criteria also required that the members slept the previous night in the household. In later surveys, this criteria was

Var Model Description

dropped and all usual residents and visitors who slept in the household the previous night were interviewed. Non *de facto* women were later dropped in the analysis and do not appear in the Individual Recode Data File.

HV011 X	Total number of eligible men indicates the number of men found eligible for the men's or husband's survey in the household. The selection criteria is country-specific and will be documented in the Household Recode Documentation for each country.
HV012	Total number of <i>de jure</i> household members gives the number of household members that usually live in the household.
HV013	Total number of <i>de facto</i> household members gives the number of household members that slept in the household the previous night, including visitors.
HV014	Number of children resident in the household and aged 5 and under. Visiting children are not included.
HV015	Result of household interview. Code 1 represents a completed interview. For all other cases, only RECH0 will exist in the data file. For flat and rectangular format data files, cases with a result code different than 1 are dropped from the file.
HV016	Day of interview.
HV017	Number of visits for the interview.
HV018	Interviewer identification code. Codes are country-specific.
HV019	Data entry keyer code. Codes are country-specific.
HV020	The ever-married sample indicator is a constant for all cases in the data file. For all woman samples it is code 0, and for ever married samples it is code 1.
HV021	Primary sampling unit is a number assigned to sample points to identify the primary sampling units for use in the calculation of sampling errors. This variable is usually the same as the cluster number and/or the ultimate area unit, but may differ if the sample design required a multistage selection process.
HV022	Sample strata defines the pairings or groupings of primary sampling units used in the calculation of sampling errors when using the Taylor series expansion method (for example, with the package Clusters).
HV023	Sample domain defines the basic geographic units within which the sample was designed. For example, if the sample was designed to be self-weighting within region, this variable would define those regions; if the sample was designed to be self-weighting within major urban areas, other urban areas and rural areas, this variable would define the major urban, other urban and rural areas. If the sample is self-weighted at the national level, this variable is code 0.
HV024	Region of residence in which the household resides. Codes are country-specific.
HV025	Type of place of residence where the household resides as either urban or rural.

<u>Var</u>	<u>Model</u>	<u>Description</u>
HV026		Size of place of residence is the type of place in which the household resides. Urban areas are classified into large cities (capital cities and cities with over 1 million population), small cities (population over 50,000), and towns (other urban areas), and all rural areas are assumed to be countryside.
HV027	X	Selection for men's or husband's survey indicates whether the household was selected for the subset of households in which the men's or husband's survey was administered. Code 1 indicates a men's survey and code 2 a husband's survey, while code 0 indicates the household was not selected.
HV028	X	Sample weight for men's or husband's survey is an 8 digit variable with 6 implied decimal places. To use the sample weight divide it by 1000000 before applying the weighting factor. All sample weights are normalized such that the weighted number of cases is identical to the unweighted number of households selected for the men's or husband's survey when using the full dataset with no other selection. This variable should be used to weight all tabulations produced using the households selected for the men's or husband's survey. For self-weighting samples this variable is equal to 1000000. For households not included in the men's or husband's survey subsample, this variable is set to zero.
HV030		Field supervisor's code. Codes are country-specific. The size of this variable was changed from 2 digits in the previous surveys to 3 digits in MEASURE DHS+.
HV031		Field editor's code. Codes are country-specific. The size of this variable was changed from 2 digits in the previous surveys to 3 digits in MEASURE DHS+.
HV032		Office editor's code. Codes are country-specific.
HV033		Ultimate area unit selection probability is the probability of selection of the ultimate area unit, ignoring the household selection. This variable can be used in conjunction with data for the sample point, such as service availability data.
HV035		Number of children under five eligible for height and weight.
HV040		Cluster altitude in meters. Used to adjust the anemia measurement for altitude.
HV041		Number of women eligible for height and weight measured.
HV042	X	Household selected for hemoglobin measurements.
HV043	X	Household selected for women's status module.
HV044	X	Household selected for the domestic violence module.

Section H1 (RECH1)

Household Schedule

<u>Var</u>	<u>Model</u>	<u>Description</u>
HVIDX		Line number of the household member.
HV101		Relationship to the head of the household.
HV102		Whether the member is a <i>de jure</i> household member, i.e., whether the member is a usual resident of the household.
HV103		Whether the member is a <i>de facto</i> household member, i.e., whether the member slept in the household the previous night.
HV104		Sex of the household member.
HV105		Age of the household member.
HV106		Highest level of education the household member attended. This is a standardized variable providing level of education in the following categories: No education, Primary, Secondary, Higher. Any member below the lower age limit for the education questions is classified in the "No education" category. Note that the lower age limit may be different from 6 years in some countries. Country-specific categorizations of education are recorded in RECH3.
HV107		Highest year of education gives the years of education completed at the level given in HV106. BASE: All household members except those answering "No education" or with missing data or the response "Don't know" for HV106 ($HV106 < 0$ & $HV106 < 9$ & $HV106 < 8$).
HV108		Education in single years. This variable is constructed from the educational level (HV106) and the grade at that level (HV107) as follows: $HV106 = > HV108$ $0 = > 0$ $1 = > HV107$ $2 = > HV107+x$ $3 = > HV107+y$ $9 = > 99$ <p>x = years to complete primary education y = years to complete primary and secondary education where both x and y are country-specific.</p>
HV109		Educational achievement recodes the education of the household member into the following categories: None, incomplete primary, complete primary, incomplete secondary, complete secondary, higher education. See related variables HV106, HV107, HV108.
HV110		Whether the household member is still in school. All members aged equal to or older than the upper limit (usually 25 years) for this question or who have not attended school are coded 0 (Not in school).
HV111		Whether the mother of the household member is still alive. BASE: All children in the household aged less than 15.

<u>Var</u>	<u>Model</u>	<u>Description</u>
HV112		Line number in the household of the mother of the member. This variable is code 00 if the mother is not a member of the household. BASE: All children in the household aged less than 15.
HV113		Whether the father of the household member is still alive. BASE: All children in the household aged less than 15.
HV114		Line number in the household of the father of the member. This variable is code 00 if the father is not a member of the household. BASE: All children in the household aged less than 15.
HV115	X	Marital status of the household member.
HV116	X	Whether the household member is currently, formerly or never married (or lived with a partner). Currently married includes married women and women living with a partner, and formerly married includes widowed, divorced, separated women and women who have lived with a partner but are not now living with a partner. In countries where the only question asked relates to whether the household member is ever married, the responses are coded 2 for ever married and 0 for never married.
HV117		Eligibility of the household member for the individual women's survey. This indicates the women included in the individual recode. In most surveys, both de facto and non de facto women are interviewed, however women are included in the individual recode only if they were eligible for interview and were de facto members of the household. (A few surveys used a de jure sample and this selection does not apply in those countries.)
HV118	X	Eligibility of the household member for the individual men's survey.
HV120		Eligibility of the child for the height/weight and hemoglobin.
HV121		Household member attended school during current school year.
HV122		Educational level attended during current school year.
HV123		Grade of education at the level of education attended during current school year.
HV124		Education in single years during current school year.
HV125		Household member attended school during previous school year.
HV126		Educational level attended during previous school year.
HV127		Grade of education at the educational level during previous school year.
HV128		Education in single years during- previous school year.
HV129		School attendance status. 0 Never attended. Children with no education. 1 Entered school. Children who did not attend school the previous year but are currently enrolled.

<u>Var</u>	<u>Model</u>	<u>Description</u>
2		Advanced. Children at a current level that is higher than the previous year
3		Repeating. Children who are at the same level than the previous year or at a level less than the previous year.
4		Dropout. Children who were at school the previous year but not currently attending school.
5		Left school 2+ years ago. Children who are not currently attending school and did not go to school the previous year.
8		Don't know

Section H2 (RECH2)

Household Characteristics

<u>Var</u>	<u>Model</u>	<u>Description</u>
HV201		Major source of drinking water for members of the household. Individual codes are country-specific, but the major categories are standard.
HV202	X	Major source of water for household use other than for drinking. Individual codes are country-specific, but the major categories are standard.
HV203	X	Whether the same source of water is used for drinking water as for household water.
HV204		Time taken to get to the water source for drinking water. BASE: All respondents except those with drinking water either piped to, or available from a well in, the residence, yard or plot or who use rainwater or bottled water (HV201 < 11 & HV201 < 21 & HV201 < 41 & HV201 < 61). The actual selection criteria is country-specific.
HV205		Type of toilet facility in the household. Individual codes are country-specific, but the major categories are standard.
		Whether the household has:
HV206		Electricity.
HV207		A radio.
HV208		A television.
HV209		A refrigerator.
		Whether any member of the household has:
HV210		A bicycle.
HV211		A motorcycle.
HV212		A car.
HV213		Main material of the floor. Individual codes are country-specific, but the major categories are standard.
HV214	X	Main material of the walls. Individual codes are country-specific, but the major categories are standard.
HV215	X	Main material of the roof. Individual codes are country-specific, but the major categories are standard.
HV216		Number of rooms used for sleeping in the household.
HV217		Relationship structure in the household describes the household composition in the following categories: one adult, two related adults of the opposite sex, two related adults of the same sex, three or more related adults, all other combinations. Only usual (de jure) members aged 15 and over are considered in determining the relationship structure.
HV218		Line number of head of household. This should always be 01, however there are some households in certain surveys in which the head of household has not been listed as the first person in the household listing.
HV219		Sex of head of household.
HV220		Age of head of household.

<u>Var</u>	<u>Model</u>	<u>Description</u>
HV221		Whether the household has a telephone.
HV222		Type of salt used for cooking in the household is used to assess the presence of iodine in the diet.
HV225		Share toilet with other households.
HV226		Type of cooking fuel.
HV227		Have a bednet for sleeping.
HV228		Children under 5 slept under bednet last night.
HV230		Place for hand washing.
HV231		Items present: Water, tap.
HV232		Items present: Soap/other cleansing agent.
HV233		Items present: Basin.
HV234		Test salt for Iodine.

Sections H3-H4 (RECH3-RECH4)

Country-Specific Household Variables

The following sections will appear in the household recode data file as needed on a country-specific basis.

- RECH3 All single occurrence country-specific variables relating to the household.
- RECH4 Country-specific variables from the household schedule. Variable IDXH4 is always included as the first variable in this section and is equal to HVIDX for each entry in the household schedule.

Sections H5 (RECH5)**Women's height/weight and hemoglobin Variables**

<u>Var</u>	<u>Model</u>	<u>Description</u>
HA0		Index to household schedule.
HA1		Women's age in years.
HA2		Respondent's weight (kilos-1d).
HA3		Respondent's height (cms-1d).
HA4		Height/Age Percentile
HA5		Height/Age Standard deviations.
HA6		Height/Age Percent ref. Median.
HA11		Weight/Height Std deviations (DHS).
HA12		Weight/Height Percent ref. median (DHS).
HA12A		Weight/Height Percent ref. median (Foggarty)
HA12B		Weight/Height Percent ref. median (WHO).
HA13		Women's result of measurement.
HA32		Date of birth (cmc).
HA33		Completeness of information.
HA35		Smoking.
HA40		Body mass index for respondent.
HA41		Rohrer's index for respondent.
HA50		Under age 18.
HA51		Line number of parent/caretaker.
HA52		Whether the consent to draw a droplet of blood was granted after reading a consent statement to woman/parent/responsible adult.
HA53		Hemoglobin level in g/dl with 1 implied decimal.
HA54		Currently pregnant.
HA55		Result of Hemoglobin measuring.
HA56		Hemoglobin level adjusted by altitude in g/dl with 1 implied decimal.
HA57		Anemia level. Levels below 7.0 g/dl are considered as severe anemia, levels between 7.1g/dl and 9.9g/dl are considered as moderate anemia and cases between 10.0 g/dl and 10.9 g/dl

are considered as mild anemia for pregnant women and between 10.0 g/dl and 11.9 g/dl for all other adult women.

HA58 X Agrees to referral. In case where the anemia level is severe, respondents are asked whether this information can be given to a doctor at a specified health facility for follow up.

Section H6 (RECH6) Children's Height/Weight/Hemoglobin Variables

<u>Var</u>	<u>Model</u>	<u>Description</u>
HC0		Index to household schedule
HC1		Age in months
HC2		Weight in kilograms (1 dec.)
HC3		Height in centimeters (1 dec.)
HC4		Height/Age Percentile
HC5		Height/Age Standard deviations
HC6		Height/Age Percent of ref. median
HC7		Weight/Age Percentile
HC8		Weight/Age Standard deviations
HC9		Weight/Age Percent of ref. median
HC10		Weight/Height Percentile
HC11		Weight/Height Standard deviations
HC12		Weight/Height Percent of ref. median
HC13		Reason not measured
HC15		Height: lying or standing
HC16		Day of birth of child
HC17		Date measured (day)
HC18		Date measured (month)
HC19		Date measured (year)
HC27		Sex of the child
HC30		Month of birth of child
HC31		Year of birth of child
HC32		Date of birth (cmc)
HC33		Completeness of information
HC51		Line number of parent/caretaker
HC52		Read consent statement

HC53	Hemoglobin level (g/dl - 1 decimal)
HC55	Result of measuring (Hemoglobin)
HC56	Hemoglobin level adjusted by altitude in g/dl with 1 implied decimal
HC57	Anemia level. Levels below 7.0 g/dl are considered as severe anemia, levels between 7.1g/dl and 9.9g/dl are considered as moderate anemia and cases between 10.0 g/dl and 10.9 g/dl are considered as mild anemia.
HC58	X Agrees to referral. In case where the anemia level is severe, respondents are asked whether this information can be given to a doctor at a specified health facility for follow up
HC60	Mother's line number from woman's questionnaire
HC61	Mother's highest educational level
HC62	Mother's highest year of education
HC63	Preceding birth interval
HC64	Birth order number

Section and Variable Description - Men

<u>Section</u>	<u>Code</u>	<u>Length</u>	<u>Class</u>	<u>Occurrences</u>		<u>Section label</u>
				<u>Min</u>	<u>Max</u>	
MREC01	M0	146	S	1	1	Respondent's Basic Data
MREC11	M1	62	S	0	1	Respondent's Basic Data
MREC22	M2	41	S	0	1	Reproduction
MREC31	M3	99	S	0	1	Contraceptive Table
MREC32	M4	295	S	0	1	Contraceptive Use
MREC51	M5	57	S	0	1	Marriage/Exposure
MREC61	M6	148	S	0	1	Fertility Preferences
MREC71	M7	46	S	0	1	Occupation and Work Status
MREC75	M8	229	S	0	1	AIDS and Condom Use
MREC91	MA	?	S	0	1	Country-specific - Single variables
MREC92	MB	?	?	0	?	Country-specific
MREC93	MC	?	?	0	?	Country-specific

? implies that the entry is country-specific

Section 01 (MREC01)

Respondent's Basic Data

Var Model Description

MCASEID	Case identification is used to uniquely identify each respondent. In most surveys this is constructed by concatenating the cluster or sample point number, the household number and the respondent's line number, but in some surveys this may be the questionnaire number taken from the front page of the questionnaire.
MV000	Alphabetic country code to identify the survey from which the data were collected. The code is based on an international standard code. This variable is 3 characters in length, with the third character indicating the format of the recode file used for this survey. For all surveys in DHS III this code will be 3. For example: BI4 is Benin, BD4 is Bangladesh, and ZW4 is Zimbabwe.
MV001	Cluster number is the number identifying the sample point as used during the fieldwork. This variable may be a composite of several variables in the questionnaire. If so, the original variables are included in MREC91 as country-specific variables.
MV002	Household number is the number identifying the household in which the respondent was interviewed, within the sample point. In some cases, this variable may be the combination of dwelling number and household number within dwelling. In these cases, the original variables are included as country-specific variables.
MV003	Respondent's line number in the household schedule.
MV004	Ultimate area unit is a number assigned to each sample point to identify the ultimate area units used in the collection of data. This variable is usually the same as the cluster number, but may be a sequentially numbered variable for samples with a more complicated structure.
MV005	Sample weight is an 8 digit variable with 6 implied decimal places. To use the sample weight divide it by 1000000 before applying the weighting factor. All sample weights are normalized such that the weighted number of cases is identical to the unweighted number of cases when using the full dataset with no selection. This variable should be used to weight all tabulations produced using the data file. For self-weighting samples this variable is equal to 1000000.
MV006	Month of interview.
MV007	Year of interview. This variable now occupies 4 digits.
MV008	Century month code of date of interview (see note on century month codes).
MV009	Month of birth of respondent (see note on imputed dates).
MV010	Year of birth of respondent (see note on imputed dates). This variable now occupies 4 digits.
MV011	Century month code of date of birth of the respondent (see note on century month codes).

<u>Var</u>	<u>Model</u>	<u>Description</u>
MV012		Current age in completed years is calculated from the century month code of the date of birth of the respondent (MV011) and the century month code of the date of interview (MV008). In a few cases the age in the data file will be different from that reported by the respondent when the respondent's birthday was in the month of interview, but he had not yet had his birthday. If the respondent correctly reported his age at his last birthday (and not his age at his next birthday) then the calculated age was rounded up from the reported age, to avoid inconsistencies between the age and the century month code for the birth.
MV013		Current age in 5-year groups is produced by grouping MV012.
MV014		Completeness of information for the date of birth of the respondent (see note on imputed dates).
MV015		Result of individual interview. Code 1 represents a completed interview. For all other cases, only MREC01 will exist in the data file. For flat and rectangular format data files, cases with a result code different than 1 are dropped from the file.
MV016		Day of the month in which the interview took place.
MV021		Primary sampling unit is a number assigned to sample points to identify the primary sampling units for use in the calculation of sampling errors. This variable is usually the same as the cluster number and/or the ultimate area unit, but may differ if the sample design required a multistage selection process.
MV022		Sample strata defines the pairings or groupings of primary sampling units used in the calculation of sampling errors when using the Taylor series expansion method''''.
MV023		Sample domain defines the basic geographic units within which the sample was designed. For example, if the sample was designed to be self-weighting within region, this variable would define those regions; if the sample was designed to be self-weighting within major urban areas, other urban areas and rural areas, this variable would define the major urban, other urban and rural areas. If the sample is self-weighted at the national level, this variable is code 0.
MV024		<i>De facto</i> region of residence. This is a copy of MV101, added to this section to allow for analysis of completion rates by region.
MV025		<i>De facto</i> type of place of residence. This is a copy of MV102, added to this section to allow for analysis of completion rates by urban/rural residence.
MV026		<i>De facto</i> place of residence is the type of place in which the respondent was interviewed. This is a copy of MV134, added to this section to allow for analysis of completion rates by type of place of residence.
MV027		Number of visits for the interview.
MV028		Interviewer identification code. Codes are country-specific.
MV029		Data entry keyer code. Codes are country-specific.

<u>Var</u>	<u>Model</u>	<u>Description</u>
MV030		Field supervisor's code. Codes are country-specific. This variable now occupies 3 digits.
MV031		Field editor's code. Codes are country-specific. This variable now occupies 3 digits.
MV032		Office editor's code. Codes are country-specific.
MV033		Ultimate area unit selection probability is the probability of selection of the ultimate area unit, ignoring the household selection. This variable can be used in conjunction with data for the sample point, such as service availability data.
MV034		Line numbers of wives as recorded in the household schedule. This is a multiple variable with entries for up to 8 wives. This variable can be used, in conjunction with the cluster or sample point number and the household number to match the men's data with the women's data, to allow for the analysis of couples. An entry with the value 0 means the wife was not listed as a member of the household.
MV034A		Men's wife or partner. For each of the women listed in MV034, it gives her status wife or partner of the men.
MV035		Number of wives or partners for whom line numbers are given in MV034.
MV801		Time of the start of the interview. The first two digits give the time in hours using the 24-hour clock, and the last two digits give the minutes within that hour.
MV802		Time of the end of interview is coded as for the start of interview.
MV803		Length of interview in minutes is calculated from the previous two variables for interviews requiring only one visit. Interviews that required more than one visit are coded 96.

Section 11 (REC11)

Respondent's Basic Data

<u>Var</u>	<u>Model</u>	<u>Description</u>
MV101		<i>De facto</i> region of residence. Region in which the respondent was interviewed. Codes are country-specific. For <i>de jure</i> region of residence, see MV139.
MV102		<i>De facto</i> type of place of residence. Type of place of residence where the respondent was interviewed as either urban or rural. Note that this is not the respondent's own categorization, but was created based on whether the cluster or sample point number is defined as urban or rural. See also MV134. For <i>de jure</i> type of place of residence, see MV140.
MV103		Childhood place of residence is classified into city, town and countryside as reported by the respondent. In some countries, additional codes are used for capital/major cities (code 0) and for abroad (code 4).
MV104		Number of years the respondent has lived in the village, town, or city where he was interviewed. Visitors to the community are coded 96.
MV105		Type of place of previous residence is coded as for MV103. In some countries, additional codes are used for capital/major cities (code 0) and for abroad (code 4). BASE: All respondents except those answering "Always" or "Visitor" to MV104 (MV104 < 95 & MV104 < 96).
MV106		Highest education level attended. This is a standardized variable providing level of education in the following categories: No education, Primary, Secondary, Higher. In some countries the educational system does not fit naturally within this scheme and a different categorization was used for the Final Report. In this case, this variable is constructed as accurately as possible from the country's own scheme and the variable used for the Final Report is included as a country-specific variable.
MV107		Highest year of education gives the years of education completed at the level given in MV106. BASE: All respondents except those answering "No education" or with missing data for MV106 (MV106 < 0 & MV106 < 9).
MV108		Literacy of the respondent. In many countries, respondents with secondary or higher levels of education are coded 1, "Reads easily." The exact criteria for this assumption is country-specific.
MV109		Whether the respondent usually reads a newspaper or magazine at least once a week.
MV110		Whether the respondent usually watches television every week.
MV111	A	Whether the respondent usually listens to a radio every day.
MV112	B	Whether the respondent usually listens to a radio every week.
MV130		Religion. Both the question and the codes are country-specific.
MV131		Ethnicity. Both the question and the codes are country-specific.

Var Model Description

MV133 Education in single years. This variable is constructed from the educational level (MV106) and the grade at that level (MV107) as follows:

MV106 =>		MV133
0	=>	0
1	=>	MV107
2	=>	MV107+x
3	=>	MV107+y
9	=>	99

x = years to complete primary education
y = years to complete primary and secondary education
where both x and y are country-specific.

MV134 *De facto* place of residence is the type of place in which the respondent was interviewed. Urban areas are classified into large cities (capital cities and cities with over 1 million population), small cities (population over 50,000), and towns (other urban areas), and all rural areas are assumed to be countryside. Note that this classification differs from that used in DHS I.

MV135 Whether the respondent is a usual resident of the household or is just visiting the household. Responses of "Visitor" to MV104 are visitors to the city, town or village where the interview took place, but MV135 shows respondents who were visitors to the household.

MV136 Total number of household members is the number of usual residents plus the number of visitors who slept in the house the previous night that were listed in the household schedule.

MV138 Number of eligible men in the household. Eligible men are usually defined to be men aged 15-59 who slept in the household the previous night, irrespective of whether they usually reside in the household or are visiting the household. In some countries an ever-married sample is used for the individual interview, and so the eligibility criteria is further restricted to husbands of eligible women.

~~MV148 Whether the respondent is still in school. In DHS III, these data are now taken directly from a question in the men's questionnaire. Men who are older than 24 years of age are coded 0, assuming that they are no longer in school.~~

MV149 Educational achievement recodes the education of the respondent into the following categories: None, incomplete primary, complete primary, incomplete secondary, complete secondary, higher education. See related variables MV106, MV107, MV133.

MV150 Relationship to the head of the household. These data are taken from the household schedule.

MV151 Sex of the head of the household.

MV152 Age of the head of the household.

Literacy

MV155 Literacy. The respondent is asked to read a written sentence and the interviewer would note whether the respondent could read it or not at all.

MV156 Ever participated in a literacy program outside of primary.

Media

<u>Var</u>	<u>Model</u>	<u>Description</u>
MV157		Frequency of reading newspaper or magazine
MV158		Frequency of listening to radio
MV159		Frequency of watching television

Travel

MV167		Times away from home in last 12 months
MV168		Away for more than one month

Tobacco smoking

MV463A		Cigarettes
MV463B		Pipe
MV463C		Other tobacco
MV463E		Country specific
MV463D		Country specific
MV463F		Country specific
MV463G		Country specific
MV463Z		Nothing
MV464		Number of cigarettes in last 24 hours. BASE: Respondents smoking cigarettes.

Section 22 (MREC22)

Reproduction

Var Model Description

MV201		Total number of children ever born.
MV202		Total number of sons living at home.
MV203		Total number of daughters living at home.
MV204		Total number of sons living away from home.
MV205		Total number of daughters living away from home.
MV206		Total number of sons who have died.
MV207		Total number of daughters who have died.
		MV201 is the sum of variables MV202 to MV207.
MV212		Age of respondent at first birth.
MV213		Whether the respondent's first listed partner is currently pregnant.
MV217		Knowledge of the women's ovulatory cycle.
MV218		Total number of living children is the sum of variables MV202 to MV205.
MV225		At the time the respondent's partner became pregnant with the current pregnancy, whether the current pregnancy was wanted then, later or not at all. BASE: Respondent's whose partner is currently pregnant (MV213 = 1).

Children's fathering

MV245		Number of women the respondent fathered the children with.
MV246		Married to the mother of the first child when he was born.

Section 31 (MREC31)

Contraceptive Table

Var Model Description

- MV301 Knowledge of any method is classified into modern, traditional and folkloric methods as follows: Modern methods are Pill, IUD, Injections, Diaphragm, Condom, Female Sterilization, Male Sterilization, Implants, Lactational amenorrhea, Female condom and Foam/Jelly. Traditional methods are Periodic Abstinence (Rhythm), Withdrawal, and Abstinence. Folkloric methods are the category "other" and any other country-specific methods. If a respondent knows both a traditional method and a modern method then the modern method takes priority and he is coded as knowing a modern method. Similarly, if a man knows a traditional method and a folkloric method, the traditional method takes priority.
- MV302 Ever use of a modern, traditional or folkloric method is created in the same way as MV301.

Contraceptive Table

The contraceptive table contains entries for 20 contraceptive methods, and for each entry gives information relating to knowledge of the method, and ever use of the method. Entries 1 to 12 are standard but entries 13 to 15 are used for country-specific methods. The methods relating to each entry are as follows:

1	Pill	11	Norplant™ or implants
2	IUD	12	Abstinence
3	Injections	13	Lactational amenorrhea
4	Diaphragm	14	Female condom
5	Condom	15	Foam or Jelly
6	Female Sterilization	16	
7	Male Sterilization	17	Country specific method 1
8	Periodic Abstinence (Rhythm)	18	Country specific method 2
9	Withdrawal	19	Country specific method 3
10	Other methods	20	Country specific method 4

For Foam/Jelly, if questions about the methods are asked separately (for example, foaming tablets in one set of questions and jelly in another set of questions), the original responses are recorded as country-specific variables and the standard variables presented in this section are a composite of the two sets of questions.

The contraceptive table contains variables MV304A to MV307 as follows:

- M304A Whether the method is modern, traditional or folkloric.
- MV304 Knowledge of the method, differentiating between spontaneous responses and probed responses for each method. If questions relating to the method were not asked in a particular country then code 8 "Not asked" is used.
- MV305 Whether the respondent has ever used the contraceptive method.
BASE: Respondents who knew of the method, either spontaneously (1) or after probing (being read a description of the method) (2) according to MV304.
- MV307 Whether the method is currently being used.

Example Contraceptive Table:

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
	Pill	IUD	Inj.	Dia.	Cond.	F.S	M.S	P.A	With	Oth.	Nor.	Abst	L.Am	F.C.	Foam	CS1	CS2	CS3	CS4	CS5
M304A	1	1	1	1	1	1	1	2	2	3	1	2	1	1	1	1	3	3	3	3
MV304	0	0	0	0	1	0	0	0	1	0	0	8	0	0	0	0	8	8	8	8
MV305					1				1											
MV307					1				0											

In this example the entries in the table are shown across the page while the variables in each entry are shown down the page. The numbers shown above the method names are the occurrence or entry number associated with that method. The respondent knew two methods, Condom and withdrawal. The respondent has used Condoms and withdrawal and currently using condoms.

Section 32 (MREC32)

Contraceptive Use

Var Model Description

MV312		Current contraceptive method.
MV312A		Most recent contraceptive method
MV312B		Contraceptive method with other woman
MV312C		Contraceptive method with other woman (2)
MV313		Type of contraceptive method categorizes the current contraceptive method as either a modern method, a traditional method, or a folkloric method.
MV313A		Most recent use by method type
MV313B		Method use by method type with other woman
MV313C		Method use by method type with other woman (2)

Pattern and intentions for future use.

Variables MV361 to MV364 relate to the respondent's past contraceptive practice and future intentions for using contraception.

MV361		Pattern of past contraceptive use. The questions relating to contraceptive use since the last birth are not asked and thus the respondent cannot be categorized as having used a method since the last birth or having only used a method before the last birth. All past users are given code 3.
MV362		Intention to use a contraceptive method in the future is based on two questions in the model questionnaires, and classifies those intending to use a method in the future by whether they intend to use that method in the next twelve months or not. The two "Unsure" categories correspond to replies of unsure about using a method in the future (unsure about use) or, for those intending to use a method in the future, unsure about whether they intend to use that method in the next twelve months (unsure about timing). In some countries, men who had never had sexual intercourse were not asked these questions, and are coded 6 on MV362. BASE: All respondents not currently using contraception (MV312 = 0).
MV363		Preferred future method for respondents intending to use a method in the future. BASE: Respondents not currently using a method, but intending to use a method in the future (MV312 = 0 & (MV362 = 1 or MV362 = 2 or MV362 = 3)).
MV364		Contraceptive use and intention shows current users of modern methods, current users of traditional methods, non-users who intend to use in the future and non-users not intending to use a method. In some countries, men who had never had sexual intercourse were not asked the questions relating to their intention to use contraception in the future, and are coded 5 on MV364.
MV366A	—	Acceptability of family planning messages being provided on radio.
MV366B	—	Acceptability of family planning messages being provided on television.
MV367		Whether the last child was wanted at that time, later or not at all. BASE: Men who have at least one child (MV201 > 0).

Reasons for Non-Use of Contraceptive Methods.

- MV375A Reason the respondent is not using a method of contraception to avoid pregnancy.
BASE: Men who are not currently using a contraceptive method and none of whose partners are currently pregnant (MV312 = 0 & MV213 <> 1).
- MV376 Reason the respondent does not intend to use a method of contraception in the future.
BASE: All men not currently using a contraceptive method and not intending to use a method in the future (MV362 = 5).

Sources of Contraception.

- Whether the respondent has heard about family planning in the last few months from any of the following sources:
- MV384A On the radio.
MV384B On the television.
MV384C In a newspaper or magazine.
~~MV384D From a poster.~~
~~MV384E From leaflets or brochures~~
- MV396 Whether a woman who is breastfeeding can get pregnant.
- MV3B0 Method known. The methods that are included are Pill, IUD and Female sterilization.
BASE: Respondents who knew the method (MV304 = 1)
- MV3B1 Method is suitable for couples planning a family

Advantages of the method

BASE: Respondents who answered to MV3B1 that the method is a suitable method (MV3B1 = 1)

- MV3B2A Simple to use
MV3B2B Effective
MV3B2C Affordable
MV3B2D No or few side effects
MV3B2E Can stop children when desired
MV3B2F No need for medical personnel
MV3B2G No risk of getting pregnant again
MV3B2H Once inserted, no daily worry
MV3B2I Country specific
MV3B2J Country specific
MV3B2K Country specific
MV3B2L Country specific
MV3B2X Other
MV3B2Z Don't know

Advantages of the method

BASE: Respondents who answered to MV3B1 that the method is not a suitable method (MV3B1 = 2)

- MV3B3A Too expensive
MV3B3B Against religion

<u>Var</u>	<u>Model</u>	<u>Description</u>
MV3B3C		May harm women's health
MV3B3D		Has side effects
MV3B3E		Increases promiscuity
MV3B3F		Can cause sterility/cannot have children
MV3B3G		Method can fail
MV3B3H		Baby in danger if pregnancy occurs
MV3B3I		Involves doctor/medical personnel
MV3B3J		Can lead to complications
MV3B3K		Country specific
MV3B3L		Country specific
MV3B3M		Country specific
MV3B3N		Country specific
MV3B3X		Other
MV3B3Z		Don't know
MV3B11		Age at first use of condom. BASE: Respondent who heard and used condoms (MV304 = 1 & MV305 = 1)
		<u>Reason of first condom use</u> BASE: Respondent who heard and used condoms (MV304 = 1 & MV305 = 1)
MV3B12A		Avoid pregnancy
MV3B12B		Avoid getting AIDS/HIV
MV3B12C		Avoid getting an STD
MV3B12D		Avoid infecting partner
MV3B12E		To experiment/try a condom
MV3B12F		Country specific
MV3B12G		Country specific
MV3B12H		Country specific
MV3B12X		Other
MV3B13		Frequency of use of condom BASE: Respondent who heard and used condoms (MV304 = 1 & MV305 = 1)
		<u>Timing of condom use</u> BASE: Respondent who heard and used condoms (MV304 = 1 & MV305 = 1)
MV3B14A		On partner's fertile days
MV3B14B		During partner's menstruation
MV3B14C		When not using some other method
MV3B14D		With a stranger
MV3B14E		With a commercial sex worker
MV3B14F		With anyone other than partner
MV3B14G		With wife/regular partner
MV3B14H		Country specific
MV3B14I		Country specific
MV3B14J		Country specific
MV3B14X		Other
		<u>Problems with condom</u>

Var Model Description

BASE: Respondent who heard and used condoms (MV304 = 1 & MV305 = 1)

MV3B15A Too expensive
 MV3B15B Embarrassing to obtain
 MV3B15C Difficult to dispose of
 MV3B15D Difficult to put on/take off
 MV3B15E Spoils the mood
 MV3B15F Diminishes pleasure
 MV3B15G Wife/partner objects, does not like
 MV3B15H Wife/partner got pregnant
 MV3B15I Inconvenient to use/messy
 MV3B15J Condom broke
 MV3B15K Country specific
 MV3B15L Country specific
 MV3B15M Country specific
 MV3B15X Other
 MV3B15Y No problem

Purchase of condoms

BASE: Respondent who used condoms at each intercourse (MV3B13 = 1)

MV3B16 Brand of condom
 MV3B17 Source of condoms
 MV3B18 Cost of condom's packet
 MV3B19 Number of condom in each packet
 MV3B20 Affordability of condoms

Other aspects of condom use

BASE: Respondent who have heard of condoms(MV304 = 1)

MV3B21A Diminish a man's sexual pleasure
 MV3B21B Inconvenient to use
 MV3B21C Can be reused
 MV3B21D Protects against disease
 MV3B21E Women has no right to tell a man whether he should use it

MV3B22 Consider getting sterilized
 BASE: Heard of male sterilization but did not use it (MV304 = 1 & MV305 = 0)

Benefits of male sterilization

BASE: Would consider sterilization or used male or female sterilization (MV3B22 =1 or MV3B22 = 5 or MV3B22 = 4)

MV3B23A Puts man in control
 MV3B23B Effective method
 MV3B23C Operation is safe
 MV3B23D Safer than female sterilization
 MV3B23E Operation is inexpensive
 MV3B23F Less expensive than female sterilization
 MV3B23G Operation is simple
 MV3B23H Gives man freedom

Var Model Description

MV3B23I Country specific
 MV3B23J Country specific
 MV3B23K Country specific
 MV3B23X Other

Reason never consider sterilization

BASE: Women who would not consider getting sterilized (MV3B22 = 2)

MV3B24A Against religion
 MV3B24B Bad for man's health
 MV3B24C Operation not safe
 MV3B24D Less intrusive ways available
 MV3B24E May want more children
 MV3B24F May remarry some day
 MV3B24G Loss of wages
 MV3B24H Loss of sexual function
 MV3B24I Loss of manliness
 MV3B24J Country specific
 MV3B24K Country specific
 MV3B24L Country specific
 MV3B24X Other

MV3B25A Contraception is woman's business and a man should not worry
 MV3B25B Sterilized women become promiscuous
 MV3B25C To a man equivalent to being castrated
 MV3B25D Woman can get pregnant not the man, so the women should be sterilized

Section 51 (MREC51)

Marriage

Var Model Description

MV501		Current marital status of the respondent.
MV502		Whether the respondent is currently, formerly or never married (or lived with a partner). Currently married includes married men and men living with a partner, and formerly married includes widowed, divorced, separated men and men who have lived with a partner but are not now living with a partner.
MV503		Whether the respondent has been married or lived with a woman once or more than once. BASE: Ever-married men (MV501 <> 0).
MV505		The number of wives the respondent currently has. This is the number of wives and live-in partners. BASE: Currently married or in union men (MV502 = 1).
MV505A		Number of wives
MV505B		Number of live-in partners

First marriage or union

Variables MV507 to MV513 relate to the date of start of the first marriage or union.

BASE: Ever-married men (V501 <> 0).

MV507		Month of start of first marriage or union (see note on imputed dates).
MV508		Year of start of first marriage or union (see note on imputed dates). This variable occupies 4 digits.
MV509		Century month code of the date of start of first marriage or union (see note on century month codes).
MV510		Completeness of information for the date of start of the first marriage or union (see note on imputed dates).
MV511		Age at start of first marriage or union is calculated from the century month code of the date of start of first marriage or union and the century month code of the date of birth of the respondent.
MV512		Years since start of first marriage or union is calculated from the century month code of the start of first marriage or union and the century month code of the date of interview.
MV513		Marital duration is actually the number of years elapsed since the start of the first marriage or union until the date of interview grouped into five-year groups, irrespective of whether the respondent is still married to his first partner.

Sexual intercourse

Variables MV525 to MV528 relate to age at first intercourse, frequency of intercourse and time since last sexual relations. BASE (for variables MV527 to MV532): Respondents who have had sexual intercourse (MV525 <> 0).

- MV525 Age at first sexual intercourse. Respondents who had never had sex are coded 0.
- MV527 Time since the last sexual relations as reported by the respondent. The first digit gives the units in which the respondent gave his answer: 1 - Days ago, 2 - Weeks ago, 3 - Months ago, 4 - Years ago,, with 9 meaning a special answer was given. The last two digits give the time in the units given. Any value for time greater than 90 is a special answer.
- MV531 Age at first sexual intercourse - imputed. This is the same as MV525, except for respondents who reported that their first sexual intercourse was at the time of their union. For these cases, the age at first sex is taken from the age at first union. In cases where the age at first sex was inconsistent with the age at conception of the first child, but only by one year (V532 = 3), the age at first sex was reduced by one year, consistent with the "Rule of one" applied in DHS I which is the correction of the data if the error is no more than a month. Other cases flagged as inconsistent on variable V532 (codes 1, 2, 4, 5) are recoded as 97 (inconsistent). Cases coded 6 on V532 are not changed.
- MV532 Flag variable for inconsistencies found in editing the responses for MV525.
- 0 No flag
 - 1 Respondent reported age at first sexual intercourse that exceeds his current age
 - 2 Respondent reported his age at first sexual intercourse as occurring more than one year after the conception of his first child
 - 3 Respondent reported his age at first sexual intercourse as occurring up to one year after the conception of his first child
 - 4 Respondent reported that his first sexual intercourse was at the time of his first marriage, but the respondent was never married
 - 5 Respondent reported that his first sexual intercourse was at the time of his first marriage, but his first marriage occurred after the conception of his first child
 - 6 Respondent reported his first sexual intercourse as being after his first marriage
- MV534 Whether the respondent, who is not currently married or living with a man, has a regular, occasional, or no sexual partner.
BASE: Men who are not currently married and not living with a woman (MV502 <> 1).
- MV534A One or more regular partner.
- MV535 Whether the respondent has ever been married or lived with a woman.
BASE: Men who are not currently married and not living with a woman (MV502 <> 1).
- MV536 Recent sexual activity. Whether the respondent had sex and whether it was during the last 4 weeks.
- MV538 Total wives, and partners including current and all former wives and partners
- MV539 Other regular sexual partners

Section 61 (MREC61)

Fertility Preferences

Var Model Description

- MV602 Fertility preferences. This variable comes primarily from a single question in the MEASURE *DHS+* questionnaires. This is the same question used in the DHS I Model "B" questionnaire, the DHS II, and the DHS III Model "A" and "B" questionnaires. However, for DHS I Model "A" questionnaires, this variable was constructed from a series of questions. Men who respond that they want another child, but when asked when they would like the next child, respond that they can not have anymore children or their wife cannot get pregnant, are classified in the "declared infecund category", and not in the "Wants another" category. These men can be identified in variable MV616, where the original response to the question asking how long they would like to wait before having another child is recorded. In some countries, men who had never had sexual intercourse were not asked the questions relating to desire for future children, and are coded 6 on MV602.
BASE: *All men.*
- MV603 Preferred waiting time before the birth of another child is created from a single question asking how long from the date of interview the respondent would like to wait before the birth of the next child. In some countries there may be some additional non-numeric responses to the question of how long to wait before the next birth. These are assigned additional codes on a country-specific basis.
BASE: *All men who want another child (MV602 = 1).*
- MV604 The preferred waiting time to the next birth is grouped into 12-month categories with responses of more than six years coded as 6+ years. Non-numeric responses are coded into one group (7 "Non-numeric"), but with "Don't know" and missing responses in their own categories (8 & 9). The additional response "Soon/Now" is not grouped with the other non-numeric codes, but is recoded as less than one year waiting time.
BASE: *All men who want another child (MV602 = 1).*
- MV605 Desire for more children is a constructed variable classifying respondents who want more children by whether they want the next child soon (less than 2 years) or they want the next child later (2+ years). In some countries, men who had never had sexual intercourse were not asked the questions relating to desire for future children, and are coded 8 on V605.
BASE: *All men.*
- MV610 Whether the respondent thinks his partner approves of couples using a method to avoid pregnancy.
BASE: *Currently married or in-union men (MV502 = 1).*
- MV611 How often the respondent discussed family planning with his partner in the past year.
BASE: *Currently married or in-union men (MV502 = 1).*
- MV612 Whether the respondent approves, in general, of couples using a method to avoid pregnancy.

<u>Var</u>	<u>Model</u>	<u>Description</u>
MV613		The ideal number of children that the respondent would have liked to have in his whole life, irrespective of the number he already has. In many countries it was possible for a respondent to reply to this question with a range of values, in which case this variable contains the midpoint between these values. If the midpoint is not an exact number then the number is rounded up in half the cases and rounded down for the other half. In situations where a range of values was collected, the original variables are included as country-specific variables. In some countries, additional country-specific categories are included, such as "It depends on God" or "As many as I can support" and are given country-specific codes.
MV614		This variable groups the preceding variable such that 6 or more children are in one category 6+ and all non-numeric responses are coded 7.
MV616		This variable records the original response to the question "How long would you like to wait from now before the birth of another child?" The first digit gives the units in which the respondent answered (1 indicates months, 2 indicates years, and 9 indicates a special response), while the last two digits give the time in those units. If the units value is 9 then the variable contains a special response, and if the duration value is greater than 90 this also indicates a special response. For example, code 994 is used for the responses "Soon/Now", and code 299 would mean that the response was given in years but the actual duration was missing on the questionnaire. BASE: All men who want another child (MV602 = 1), plus those originally responding that they want another child, but then say their partner cannot get pregnant (see also MV602).
MV621		Whether the respondent believes his partner wants the <u>same</u> number of children, <u>more</u> children or <u>fewer</u> children than he wants herself. BASE: Currently married or in union men (MV502 = 1).
MV627		Ideal number of boys.
MV628		Ideal number of girls.
MV629		Ideal number of either sex. These three variables should sum to the total ideal number of children given in variables MV613. If the response to the question for variables MV613 is a non-numeric response, these variables are coded with the same response. In addition, there may be non-numeric responses on each of these questions. Country specific categories for non-numeric responses may also be recorded for these variables.

<u>Var</u>	<u>Model</u>	<u>Description</u>
		Whether the respondent discussed the practice of family planning with any of the following people:
MV630A		Husband or partner
MV630B		Mother
MV630C		Father
MV630D		Sister(s)
MV630E		Brother(s)
MV630F		Daughter(s)
MV630G		Mother-in-law
MV630H		Friends or neighbors
MV630I		<i>Country specific</i>
MV630J		<i>Country specific</i>
MV630K		<i>Country specific</i>
MV630L		<i>Country specific</i>
<i>MV630M</i>		<i>Country specific</i>
<i>MV630N</i>		<i>Country specific</i>
<i>MV630O</i>		<i>Country specific</i>
MV630X		Other people
MV631		Problem if became pregnant BASE: Wife listed first is not currently pregnant (MV642(1) <> 1).
		<u>Reason for not having sex</u>
MV633A		Husband has STD
MV633B		Husband has other women
MV633C		Recent birth
MV633D		Tired, mood
MV633E		<i>Country specific</i>
MV633F		<i>Country specific</i>
MV633G		<i>Country specific</i>
		<u>Husband's rights</u>
MV634A		Get angry
MV634B		Refuse financial support
MV634C		Use force for unwanted sex
MV634D		Have sex with another women
MV635		Wife number would prefer next child with BASE: Respondent with more than one partner who wants to have another child (MV602 = 1 & MV035 <> 0).
MV636		Planning to have children with any other partner
MV637		Plan to take another wife
MV638		Main reason may take another wife
MV640		Sequence number of partners
MV641		Line number of wives/partners

<u>Var</u>	<u>Model</u>	<u>Description</u>
MV642		Partner currently pregnant
MV643		Current pregnancy wanted
MV644		Problem if became pregnant
MV645		Wives' desire for children
MV646		Discussed FP with partner
MV647		Wife approves FP
MV648		Discussed FP with health worker
MV649A		Interest in avoiding unwanted pregnancies
MV649B		Interest in helping his partner to have a safe pregnancy.
MV649C		Interest in helping to care for newborn infants

Section 71 (MREC71)

Occupation and Work Status

<u>Var</u>	<u>Model</u>	<u>Description</u>
MV714		Whether the respondent is currently working.
MV716		Respondent's occupation as collected in the country. Codes are country-specific. BASE: Men who are currently working or who have worked in the last 12 months (MV731 = 1 or MV731 = 2).
MV717		Standardized respondent's occupation groups. Agricultural categories also include fishermen, foresters and hunters and are <u>not</u> the basis for selection of agricultural/non-agricultural workers. In countries, where it is not possible to differentiate between self-employed agricultural workers and agricultural employees, no attempt has been made to use other information, and code 4 has been used for both categories. The analyst may wish to use other related information to differentiate between these two categories. BASE: Men who are currently working or who have worked in the last 12 months (MV731 = 1 or MV731 = 2).
MV719		Whether the respondent works for a family member, for someone else or is self-employed. BASE: Men currently working (MV731 = 1 or MV731 = 2).
MV721	X	Works at home or away.
MV731		Whether the respondent worked in the last 12 months.
MV732		Whether the respondent works throughout they year, seasonally, or just occasionally. BASE: Men who are currently working or who have worked in the past year (MV731 = 1 or MV731 = 2).
MV733		For seasonal or part year workers, the number of months they worked in the last twelve months. BASE: Men who are working seasonally or for part of the year (MV732 = 2).
MV736		Usual amount the respondent earns in cash for the work he does. This variable is 8 digits in size. The first digit gives the units in which the amount was specified, while the remaining digits give the total amount. The first digit, or units digit, is coded as follows: 1 — per hour 2 — per day 3 — per week 4 — per month 5 — per year For example, 30000400 would indicate that the respondent received 400 per week. The currency and units of currency used are country specific. BASE: Men paid cash for their work (MV7?? = 1).
V740		Whether the respondent works on his own land, family land, rented land or on someone else's land. BASE: Men who are currently working or who have worked in the last 12 months, and who work or worked in agriculture (MV716 = country-specific agricultural category).

MV741 Type of earnings for work

MV742 Portion of the household expenditures the respondent's earnings pay.
BASE: Respondents working for cash (MV741 = 1 or MV741 = 2).

Final say of the respondent on key issues

MV743A Own health care
MV743B Making large household purchases
MV743C Making household purchases for daily needs
MV743D Visits to family or relatives
MV743E Food to be cooked each day
MV743F Deciding what to do with money wife earns
MV743G Deciding how many children to have

Justification of wife's beating

MV744A Goes out without telling him
MV744B Neglects the children
MV744C Argues with him
MV744D Refuses to have sex with him
MV744E Burns the food

MV745 Activity last 12 months.
BASE: Respondents who worked last 12 months (MV714 = 1 or MV716 = 1).

Section 75 (MREC75)

AIDS and Condom Use

Var Model Description

MV750 Heard of sexually transmitted disease
 MV751 Whether the respondent has ever heard of AIDS (Acquired Immune Deficiency Syndrome).

~~Sources of information from which the respondent has learned most about AIDS.~~

- ~~MV752A Radio~~
- ~~MV752B Television~~
- ~~MV752C Newspapers or magazines~~
- ~~MV752D Pamphlets or posters~~
- ~~MV752E Clinics or health workers~~
- ~~MV752F Churches or mosques~~
- ~~MV752G Schools or teachers~~
- ~~MV752H Community meetings~~
- ~~MV752I Friends or relatives~~
- ~~MV752J Work place~~
- ~~MV752K Country specific~~
- ~~MV752L Country specific~~
- ~~MV752M Country specific~~
- ~~MV752N Country specific~~
- ~~MV752O Country specific~~
- ~~MV752X Other responses~~

~~BASE: Men who have heard of AIDS (MV751 = 1).~~

MV753 Whether the respondent believes there is anything a person can do to avoid AIDS.

BASE: Men who have heard of AIDS (MV751 = 1).

Ways in which the respondent thinks people can avoid AIDS

BASE: Men who have heard of AIDS (MV751 = 1).

- MV754A "Safe Sex"
- MV754B Abstaining from sex
- MV754C Using condoms during sex
- MV754D Having only one sexual partner
- MV754E Avoiding sex wit prostitutes
- MV754F Avoiding sex with homosexuals
- MV754G Avoiding blood transfusions
- MV754H Avoiding injections
- MV754I Avoiding kissing
- MV754J Avoiding mosquito bites
- MV754K Seeking protection from a traditional healer
- MV754L Country specific
- MV754M Country specific
- MV754N Country specific
- MV754O Country specific
- MV754P Country specific
- MV754Q Country specific
- MV754R Country specific
- MV754S Country specific

MV754T *Country specific*
 MV754U *Country specific*
 MV754V *Country specific*
 MV754X Other responses
 MV754Z Does not know any means of avoiding AIDS

Ways to reduce AIDS

BASE: Respondents who have heard of AIDS (MV751 = 1).

MV754CP Reduce chances of AIDS by always using condoms during sex
 MV754DP Reduce chance of AIDS: have 1 sex partner with no other partner

Means of getting AIDS

BASE: Respondents who have heard of AIDS (MV751 = 1).

MV754JP Get AIDS from mosquito bites
 MV754WP Get AIDS by sharing food with person who has AIDS

What "Safe Sex" means to the respondent

~~MV755B — Abstaining from sex~~
~~MV755C — Using condoms during sex~~
~~MV755D — Having only one sex partner~~
~~MV755E — Avoiding sex with prostitutes~~
~~MV755F — Avoiding sex with homosexuals~~
~~MV755L — Country specific~~
~~MV755M — Country specific~~
~~MV755N — Country specific~~
~~MV755O — Country specific~~
~~MV755X — Other responses~~
~~MV755Z — Does not know the meaning of "Safe sex"~~

BASE: Men who responded that "Safe sex" was a way of avoiding AIDS (MV754A = 1).

MV756 Whether the respondent believes it is possible for a healthy-looking person to have the AIDS virus.
 BASE: Men who have heard of AIDS (MV751 = 1).

~~MV757 — Whether the respondent believes that AIDS is a fatal disease.~~
~~BASE: Men who have heard of AIDS (MV751 = 1).~~

~~MV758 — Whether the respondent believes his risk of getting AIDS is small, moderate, great, no risk at all, or that he already has AIDS.~~
~~BASE: Men who have heard of AIDS (MV751 = 1).~~

Acceptability of messages on AIDS in the media

MV759A Radio
 MV759B Television
 MV759C Newspapers
 MV759D Country specific
 MV759E Country specific

Var Model Description

MV759F Country specific
 MV759G Country specific
 MV759H Country specific
 MV759I Country specific

Ways in which the respondent has changed his sexual behavior, since hearing about AIDS, in order to avoid getting AIDS:

MV760A Did not start sex
 MV760B Stopped all sex
 MV760C Started using condoms during sex
 MV760D Restricted the number of partners to one
 MV760E Reduced the number of partners
 MV760F Ask spouse to be faithful
 MV760G No more homosexual contacts
 MV760I Stopped receiving injections
 MV760L *Country specific*
 MV760M *Country specific*
 MV760N *Country specific*
 MV760O *Country specific*
 MV760P Ask spouse to avoid prostitutes
 MV760V No non-sexual change in behavior
 MV760W Other (non-sexual) responses
 MV760X Other (sexual) responses
 MV760Y Did not change behavior
 MV760Z Don't know whether they changed behavior

BASE: Men who have heard of AIDS and have ever had sexual intercourse (MV751 = 1 & MV525 > 0).

MV761 Whether the respondent used a condom the last time he had sexual intercourse.
 BASE: Men who have ever had sexual intercourse (MV525 > 0).

Condom use during sexual intercourses

MV761B With other woman (1)
 MV761C With other woman (2)
 MV761D For money

MV762 Source of condoms known by the respondent. If the respondent does not know where to get condoms, this variable is coded 98.

BASE: Men who have ever had sexual intercourse (MV525 > 0).

Source for male condom

MV762AA Government hospital
 MV762AB Government health center or post
 MV762AC Family planning clinic
 MV762AD Mobile clinic
 MV762AE Public field worker
 MV762A Other public
 MV762AG Public country specific
 MV762AH Public country specific

<u>Var</u>	<u>Model</u>	<u>Description</u>
MV762AI		Public country specific
MV762AJ		Private hospital, clinic
MV762AK		Pharmacy
MV762AL		Private doctor
MV762AM		Private mobile clinic
MV762AN		Private field worker
MV762AO		Other private
MV762AP		Private country specific
MV762AQ		Private country specific
MV762AR		Private country specific
MV762AS		Shop
MV762AT		Church
MV762AU		Friends, relatives
MV762AV		Country specific
MV762AW		Country specific
MV762AX		Other
MV762AZ		Don't know source for condom

Source for female condoms

MV762BA		Government hospital
MV762BB		Government health center/post
MV762BC		Family planning clinic
MV762BD		Mobile clinic
MV762BE		Public field worker
MV762BF		Other public
MV762BG		Public country specific
MV762BH		Public country specific
MV762BI		Public country specific
MV762BJ		Private hospital, clinic
MV762BK		Pharmacy
MV762BL		Private doctor
MV762BM		Private mobile clinic
MV762BN		Private field worker
MV762BO		Other private
MV762BP		Private country specific
MV762BQ		Private country specific
MV762BR		Private country specific
MV762BS		Shop
MV762BT		Church
MV762BU		Friends, relatives
MV762BV		Country specific
MV762BW		Country specific
MV762BX		Other
MV762BZ		Don't know source for female condom

~~MV764 — Whether the respondent has ever heard of condoms for contraceptive use or for use to prevent STDs.~~

~~MV763 — Partners in the last 12 months.~~

Var Model Description

Sexually transmitted diseases last 12 months

MV763A Any STD
 MV763B Genital sore or ulcer
 MV763C Genital discharge
 MV763D Country specific
 MV763E Country specific
 MV763F Country specific
 MV763G Country specific

~~MV765 Whether the respondent has ever used condoms for contraceptive use or for use to prevent STDs.
 Both of these variables are created from responses to several questions in the questionnaire, but the set of questions used may vary from country to country, depending on the depth of questioning used in a particular country.~~

Reason used condom during the last sexual intercourse

MV765A Last intercourse
 MV765B With other woman (1)
 MV765C With other woman (2)
 BASE: Respondents who had sex during the last 12 months.

MV766A Number of women other than the wife the respondent had sex with in last 12 months.
 MV766B Number of women the respondent had sex with including his wife in last 12 months.

Relationship with last sexual partner

MV767A Last partner
 MV767B Other partner (1)
 MV767C Other partner (2)
 BASE: Respondents who had sexual intercourse last 12 months.

Length of time knows last sexual partner

MV768A Last partner
 MV768B Other partner (1)
 MV768C Other partner (2)
 BASE: Respondents who had sexual intercourse last 12 months.

MV769 Could get a male condom

MV769A Could get a female condom

MV770 Seek advice for last disease.
 BASE: Respondents who had an infection in the last 12 months.

Var Model Description

Place where sought advice for last disease

BASE: Respondents who sought advice for the last disease (MV770 = 1).

MV770A Clinic, hospital, private doctor
 MV770B Traditional healer
 MV770C Shop, pharmacy
 MV770D Friends, relatives
 MV770E Country specific
 MV770F Country specific
 MV770G Country specific
 MV770H Country specific
 MV770I Country specific

MV771 Advise partner when had disease.
 BASE: Respondents who had an infection in the last 12 months.
 MV772 Tried to avoid infecting partner.
 BASE: Respondents who had an infection in the last 12 months.

What was done to avoid infecting the partner

MV773A No sexual intercourse
 MV773B Used condoms
 MV773C Took medicines
 MV773D Country specific
 MV773E Country specific
 MV773F Country specific
 MV773X Other
 BASE: Respondents who tried to avoid infecting their partners (MV772 = 1).

MV774 AIDS transmitted from mother to child

Mother to child AIDS transmission knowledge

MV774A During pregnancy
 MV774B During delivery
 MV774C By breastfeeding.
 BASE: Respondents who said that the AIDS virus can be transmitted from the mother to the child (MV774 = 1).

MV775 Knows someone who has or died of AIDS.
 BASE: Respondents who have heard of AIDS (MV751 = 1).

MV776 Spoken with spouse about avoiding AIDS.
 BASE: Respondents in union or living with a man (MV502 = 1) and have heard of AIDS (MV751 = 1).

MV77 Allowed to keep AIDS infection secret.
 BASE: Respondents who have heard of AIDS (MV751 = 1).

Var Model Description

MV778 Willing to care for relative with AIDS.
BASE: Respondents who have heard of AIDS (MV751 = 1).

MV779 Person with AIDS allowed to continue teaching.
BASE: Respondents who have heard of AIDS (MV751 = 1).

MV780 Should children be taught about condoms.
BASE: Respondents who have heard of AIDS (MV751 = 1).

AIDS tests

MV781 Ever been tested
MV782 Want to be tested
MV783 Know a place to get AIDS test
MV784 Where to get an AIDS test

MV785 Heard about other STDs

Man's symptoms of sexually transmitted infections (STIs)

MV785A Abdominal Pain
MV785B Genital discharge or dripping
MV785C Foul smelling discharge
MV785D Burning pain on urination
MV785E Redness, inflammation of the genital area
MV785F Swelling in the genital area
MV785G Genital sores or ulcers
MV785H Genital warts
MV785I Genital itching
MV785J Blood in urine
MV785K Loss of weight
MV785L Impotence
MV785M Country specific
MV785N Country specific
MV785O Country specific
MV785P Country specific
MV785Q Country specific
MV785X Other
MV785Y No symptoms
MV785Z Don't know
BASE: Men who have heard about other STDs (MV785 = 1).

Woman's symptoms of sexually transmitted infections (STIs)

MV786A Abdominal pain
MV786B Genital discharge
MV786C Foul smelling discharge
MV786D Burning pain on urination
MV786E Redness or inflammation of the genital area
MV786F Swelling in the genital area
MV786G Genital sores or ulcers

<u>Var</u>	<u>Model</u>	<u>Description</u>
MV786H		Genital warts
MV786I		Genital itching
MV786J		Blood in urine
MV786K		Loss of weight
MV786L		Hard to get pregnant or to have a child
MV786M		Country specific
MV786N		Country specific
MV786O		Country specific
MV786P		Country specific
MV786Q		Country specific
MV786X		Other
MV786Y		No symptoms
MV786Z		Don't know
		BASE: Men who have heard about other STDs (MV785 = 1).

Alcohol drinking during the last 3 months

MV787 X	Number of days drank alcohol
MV788 X	Number of days gotten drunk

Injections received last 3 months

MV789 X	Number of injections in last 3 months
MV790 X	Person who gave last injection

Paid sex

MV791	Ever paid for sex
MV792	Time since last paid for sex.

Var Model DescriptionLast Sexual Intercourse, Condom Use and Other Partners

Variables MV850 to MV852 relate to the time since last sexual intercourse and whether condoms were used at that time, for sexual relationships with the respondent's husband or partner and with other people, respectively. Variable MV852 gives the number of other partners.

BASE (for MV850 to MV852): Currently married or in union men (MV502 = 1)

- MV850 — Time since the respondent's last sexual intercourse with his partner. The first digit gives the units in which the respondent gave his answer: 1—Days ago, 2—Weeks ago, 3—Months ago, 4—Years ago, with 9 meaning a special answer was given. The last two digits give the time in the units given. Any value for time greater than 90 is a special answer.
- MV850A — Whether a condom was used the last time the respondent had sexual intercourse with his partner.
- MV851 — Time since the last sexual intercourse with someone other than the respondent's partner. Respondents who have had no other partner in the prior twelve months are coded 995.
- MV852 — Whether a condom was used the last time the respondent had sexual intercourse with someone other than his partner.
- MV853 — Number of partners other than the wife or partner with whom the respondent lives, with whom the respondent had sexual intercourse in the 12 months prior to the interview.

Knowledge of Sexually Transmitted Diseases (STDs)

Variables MV855 to MV856Z relate to knowledge of sexually transmitted diseases.

BASE (for MV856A to MV856Z): Heard of any sexually transmitted disease (MV855 = 1).

- MV855 — Whether the respondent has ever heard of any sexually transmitted diseases.
- MV856A — Knowledge of Syphilis
- MV856B — Knowledge of Gonorrhea
- MV856C — Knowledge of AIDS
- MV856D — Knowledge of Genital warts
- MV856E — Knowledge of country specific
- MV856F — Knowledge of country specific
- MV856G — Knowledge of country specific
- MV856H — Knowledge of country specific
- MV856I — Knowledge of country specific
- MV856J — Knowledge of country specific
- MV856K — Knowledge of country specific
- MV856X — Knowledge of other diseases
- MV856Z — Does not know sexually transmitted diseases by name.

Var Model Description

Prevalence of STDs

Variables MV857 to MV858Z relate to whether the respondent has had any sexually transmitted disease in the twelve months prior to the interview.

BASE (for MV858A to MV858Z): Whether the respondent has ever had a sexually transmitted disease in the twelve months prior to the interview (MV857 = 1).

~~MV857 Whether the respondent has had a sexually transmitted disease in the twelve months prior to the interview.~~

~~BASE: Men who had ever had sexual intercourse and had heard of sexually transmitted diseases (MV525 <= 0 & MV855 = 1).~~

~~MV858A Last 12 months had syphilis~~
~~MV858B Last 12 months had gonorrhea~~
~~MV858C Last 12 months had AIDS~~
~~MV858D Last 12 months had genital warts~~
~~MV858E Last 12 months had *country specific*~~
~~MV858F Last 12 months had *country specific*~~
~~MV858G Last 12 months had *country specific*~~
~~MV858H Last 12 months had *country specific*~~
~~MV858I Last 12 months had *country specific*~~
~~MV858J Last 12 months had *country specific*~~
~~MV858X Last 12 months had other disease~~
~~MV858Z Last 12 months had unknown disease~~

Treatment of STDs

Variables MV859 to MV860Z relate to treatment of sexually transmitted diseases (STDs).

BASE (for MV860A to MV860Z): Whether sought advise for the last sexually transmitted disease (MV859 = 1).

~~MV859 ought advise for the last sexually transmitted disease.~~

~~BASE: Men who had a sexually transmitted disease in the 12 months prior to interview (MV857 = 1).~~

~~MV860A Sought advice from a government hospital~~
~~MV860B Sought advice from a government health center~~
~~MV860C Sought advice from a health post or dispensary~~
~~MV860D Sought advice from a mobile clinic~~
~~MV860E Sought advice from a community health worker~~
~~MV860F Sought advice from a family planning clinic~~
~~MV860G Sought advice from *country specific* public sector~~
~~MV860H Sought advice from *country specific* public sector~~
~~MV860I Sought advice from other public sector sources~~
~~MV860J Sought advice from a private hospital or clinic~~
~~MV860K Sought advice from a private pharmacy~~
~~MV860L Sought advice from a private doctor~~
~~MV860M Sought advice from a private mobile clinic~~
~~MV860N Sought advice from a community health worker~~
~~MV860O Sought advice from *country specific* medical private sector~~

Var Model Description

MV860P	—	Sought advice from <i>country specific</i> medical private sector
MV860Q	—	Sought advice from <i>country specific</i> medical private sector
MV860R	—	Sought advice from other private sector sources
MV860S	—	Sought advice from a shop
MV860T	—	Sought advice from a traditional practitioner
MV860U	—	Sought advice from relatives or friends
MV860V	—	Sought advice from <i>country specific</i> other sector
MV860W	—	Sought advice from <i>country specific</i> other sector
MV860X	—	Sought advice from other sources
MV860Z	—	Sought advice from an unknown source

Avoidance of infecting others with STD

Variables MV861 to MV863X provide information on what the respondent did when he realized that he had a sexually transmitted disease.

BASE (for MV863A to MV863X): Whether the respondent tried to avoid infecting his partner (MV863 = 1).

MV861 — Whether the respondent advised his partner when he had the sexually transmitted disease.
BASE: Men who had a sexually transmitted disease in the twelve months prior to the survey (MV857 = 1).

MV862 — Whether the respondent tried to avoid infecting his partner.
BASE: Men who had a sexually transmitted disease in the twelve months prior to the survey (MV857 = 1).

Means of avoiding infecting his partner:

MV863A	—	No sexual intercourse
MV863B	—	Used condoms
MV863C	—	Took medicines
MV863D	—	<i>Country specific</i>
MV863E	—	<i>Country specific</i>
MV863F	—	<i>Country specific</i>
MV863W	—	Other (non-sexual)
MV863X	—	Other (sexual)

Knowledge of Means of Transmission of AIDS

Variables MV864A to MV864Z provide information on the ways in which the respondent believes a person can get AIDS. Variables MV865 to MV867 relate specifically to whether AIDS can be cured, whether it is transmitted from mother to child, and whether the respondent knows anyone who has AIDS or who has died of AIDS.

BASE: Respondents who have heard of AIDS (MV751 = 1).

MV864A	—	Get AIDS from sexual intercourse
MV864B	—	Get AIDS from sex with multiple partners
MV864C	—	Get AIDS from sex with prostitutes
MV864D	—	Get AIDS from not using a condom
MV864E	—	Get AIDS from homosexual contact
MV864F	—	Get AIDS from blood transfusions
MV864G	—	Get AIDS from injections

Var Model Description

MV864H — Get AIDS from kissing
MV864I — Get AIDS from mosquito bites
MV864J — Get AIDS from *country specific*
MV864K — Get AIDS from *country specific*
MV864L — Get AIDS from *country specific*
MV864M — Get AIDS from *country specific*
MV864N — Get AIDS from *country specific*
MV864O — Get AIDS from *country specific*
MV864P — Get AIDS from *country specific*
MV864X — Get AIDS from: other responses
MV864Z — Get AIDS from unknown sources

MV865 — Whether the respondent believes AIDS can be cured.
MV866 — Whether the respondent believes that AIDS can be transmitted from mother to child.
MV867 — Whether the respondent knows someone who has AIDS or who has died of AIDS.

Reasons Respondent Assesses Risk of Getting AIDS to be Low

Variables MV868B to MV868Z give the reasons the respondent believes that he is at no risk or at a small risk of getting AIDS:

BASE: Respondents reporting they are at no risk or at only a small risk of getting AIDS (MV758 = 0 or MV758 = 1):

MV868B — No/small risk: abstains from sex
MV868C — No/small risk: uses condoms
MV868D — No/small risk: has only one sexual partner
MV868E — No/small risk: has a limited number of partners
MV868F — No/small risk: spouse has no other partners
MV868G — No/small risk: has no homosexual contact
MV868H — No/small risk: has not received a blood transfusion
MV868I — No/small risk: has had no injections
MV868J — No/small risk: country specific
MV868K — No/small risk: country specific
MV868L — No/small risk: country specific
MV868P — No/small risk: avoids prostitutes
MV868X — No/small risk: other reasons
MV868Z — No/small risk: reasons unknown

Reasons Respondent Assesses Risk of Getting AIDS to be High

Variables MV869C to MV869Z give the reasons the respondent believes that he is at a moderate or great risk of getting AIDS:

BASE: Respondents reporting they are at a moderate or great risk of getting AIDS (MV758 = 2 or MV758 = 3):

MV869C — Great/moderate risk: not using condoms
MV869D — Great/moderate risk: has more than one sex partner
MV869E — Great/moderate risk: has many sex partners
MV869F — Great/moderate risk: spouse has other sexual partners
MV869G — Great/moderate risk: has homosexual contacts
MV869H — Great/moderate risk: has received blood transfusions

Var Model Description

MV869I ——— Great/moderate risk: has had injections
MV869J ——— Great/moderate risk: country specific
MV869K ——— Great/moderate risk: country specific
MV869L ——— Great/moderate risk: country specific
MV869P ——— Great/moderate risk: frequents prostitutes
MV869X ——— Great/moderate risk: other reasons
MV869Z ——— Great/moderate risk: reasons unknown

Condom Use in Relation to AIDS

MV870 ——— Whether the respondent has heard of using condoms to avoid AIDS.

MV871 ——— Whether the respondent has ever used a condom to avoid AIDS.
BASE: Men who reported having heard of using condoms to avoid AIDS and who had ever had sexual intercourse (MV870 = 1 & MV525 \diamond 0).

Payments or Gifts for Sexual Intercourse

MV872 ——— Whether the respondent ever received or gave money or gifts in return for sexual intercourse.
BASE: Men who had ever had sexual intercourse (MV525 \diamond 0).

Prevalence of STDs

MV873 ——— Whether the respondent has suffered from a discharge from his penis during the last 12 months.
BASE: Men who had ever had sexual intercourse and had heard of sexually transmitted diseases (MV525 \diamond 0 & MV855 = 1).

MV874 ——— Whether the respondent had a sore or ulcer on his penis in the last 12 months.
BASE: Men who had ever had sexual intercourse and had heard of sexually transmitted diseases (MV525 \diamond 0 & MV855 = 1).

Last Sexual Intercourse

MV875 ——— Whether the last sexual partner was the respondent's wife or woman that he lives with, regular partner, acquaintance, someone paid for sex or someone else.
BASE: Men who had ever had sexual intercourse (MV525 \diamond 0).

Sections 91-93 (MREC91-MREC93)

Country-Specific Variables

The following sections will appear in the recode data file as needed on a country-specific basis.

- MREC91 All single occurrence country-specific variables relating to the respondent.
- MREC92-93 The last two country-specific sections are not assigned to any particular section of the questionnaire, but are used for additional modules not usually incorporated in the questionnaires.

Dictionary Listing

The dictionary listing provides the basic information relating to each variable in the data file. The dictionary listing provided separately is for rectangular and hierarchical data files. It contains a description of all of the standard variables included in the recode file. The first page gives dictionary information about the file, including the name of the dictionary, its creation date and last modification date, the questionnaire identification fields and the section identification fields. This is followed by the section descriptions giving the following information:

Section name	Name by which the section is referred.
Code	Code used to identify the record for this section.
Length	Number of characters used in the record.
Class	Whether the section is a single (S) or multiple (M) section.
Occurs	Minimum and maximum number of occurrences allowed for the section. If the maximum is greater than one then the section is a multiple section, but if the maximum is one then the section is a single section. If the minimum number of occurrences is zero then the section is not always required for every case.
Group	Maximum number of occurrences of a group within a single section, the starting location of the group within the section and the total length of all of the variables in one occurrence of the group.
Section label	Title for the section.

This is followed by the detailed description of each variable in the data file, section by section. The following information is provided for each variable:

Variable name	Name by which the variable is referred.
Location	Character position on the record.
Length	Size of the variable in characters.
Decimals	Number of decimal places in the variable. If decimal places are specified then the variable is stored with the decimal point in the data file. For example, if a variable is 4 characters in size, with 2 decimal places the variable will appear as X.XX in the data file.
Format	N is for numeric, A for alphabetic.
Class	S is for single variables in single or multiple sections, M for multiple variables of a group in single sections.
Variable label	Title of the variable.
Value labels	Labels assigned to each code for the variable.
Ranges	Pairs of values giving the lower and upper limits for the values of the variable.

The dictionary listing following is for the hierarchical data structure. The rectangular data structure has exactly the same format, but with the minimum number of occurrence of each section equal to the maximum number of occurrences of the section. The flat file data structure contains the same variables, but with all variables on one record. The locations of each variable can be calculated by concatenating all of the records end to end, but leaving out the section identification from all records and the respondent identification from all sections except the first.