

KENYA

2022 Demographic and Health Survey

HEALTH AND WELL-BEING

Maternal Health


2 in 3 women had 4 or more ANC visits during pregnancy.

89% of births were delivered by a skilled provider. *

* Skilled provider is a doctor, nurse, midwife, or clinical officer.

Number of antenatal care (ANC) visits

Percentage of women age 15-49 who had a live birth in the 2 years before the survey who had 4+ ANC visits, by residence



Malaria

Access and use of insecticide-treated nets (ITNs) is higher in malaria endemic areas.

50% of the population has access to ITNs.


43% of the population slept under an ITN the night before the survey.



Photo by Thomas Orindi/Department for International Development

Access to and use of insecticide-treated nets (ITNs)

Percentage of population with access to an ITN and percentage of population who slept under an ITN the night before the survey, by endemicity zone



Child Health

4 in 5 children age 12-23 months have been fully vaccinated against tuberculosis, polio, tetanus, pertussis, diphtheria, and measles.




80% of girls and **76%** of boys age 24-59 months are developmentally on-track in health, learning, and psychosocial well-being.

Nutritional status of children

Child nutrition has improved in the last two decades. The proportion of children under age 5 who are stunted (too short for their age) has halved, from **36%** to **18%**.

Trends in child growth measures

Percentage of children under age 5 who are malnourished



Note: Data from 2003 and later are nationally representative, while data collected before 2003 exclude the North Eastern region and several northern districts in the Eastern and Rift Valley regions.

Risk Factors and Screening for Chronic Conditions

Breast and cervical cancer screening is low. **14%** of women have ever been screened for breast cancer and **17%** for cervical cancer.

Alcohol and tobacco use

Percentage of women and men age 15-49 who consumed at least one alcoholic drink in the last month or use any type of tobacco

■ Women ■ Men

Tobacco and alcohol use is higher in men than in women.

12% of men use tobacco and **26%** use alcohol.

