


ERRATA

National Statistical Office (NSO) [Malawi], and ORC Macro. 2005. *Malawi Demographic and Health Survey 2004*. Calverton, Maryland: NSO and ORC Macro.

Page 172, first paragraph, line 13 – Text should read “. . . vitamin A in the 24 hours preceding . . .”

Page 177, first paragraph, lines 9 and 10 – Text should read “. . . 44 percent of women are anaemic; 32 percent have mild anaemia . . .”

Page 246 – Figure 13.2 is incorrect. The correct Figure 13.2 is below.



Page 261 – Table 14.7 is incorrect. The correct Table 14.7 is below.

Table 14.7 Prevalence and prompt treatment of fever by background characteristics					
Percentage of children under five years of age with fever in the two weeks preceding the survey, and among children with fever, percentage who received antimalarial drugs and who received the drugs the same/next day after developing the fever, by background characteristics, Malawi 2004					
Background characteristics	Percentage of children with fever	Number of children	Among children with fever:		
			Percentage who received antimalarial drugs	Percentage who received antimalarial drugs same/next day	Number of children
Age in months					
< 6	30.7	1,109	14.3	12.6	341
6-11	53.2	1,188	29.7	22.0	632
12-23	49.5	2,194	31.4	25.2	1,087
24-35	39.5	1,743	31.1	26.4	689
36-47	28.8	1,741	26.2	20.1	501
48-59	21.1	1,802	28.6	22.9	381
Sex					
Male	37.1	4,839	28.7	23.0	1,793
Female	37.2	4,938	28.2	22.5	1,837
Residence					
Urban	29.9	1,341	42.0	36.5	401
Rural	38.3	8,436	26.7	21.0	3,230
Region					
Northern	28.4	1,239	38.2	30.0	351
Central	39.9	4,071	24.5	20.3	1,624
Southern	37.1	4,468	30.2	23.6	1,655
District					
Blantyre	29.4	670	31.7	27.8	197
Kasungu	40.0	471	21.8	15.0	188
Machinga	35.6	405	30.1	21.3	144
Mangochi	36.8	566	24.7	17.7	208
Mzimba	28.9	630	31.0	22.5	182
Salima	42.1	281	32.0	22.7	118
Thyolo	47.3	514	40.0	27.0	243
Zomba	40.1	498	38.5	26.3	200
Lilongwe	38.3	1,376	27.2	25.0	528
Mulanje	44.3	375	29.0	25.4	166
Other districts	36.5	3,992	25.7	21.6	1,455
Education					
No education	37.3	2,594	21.2	16.3	967
Primary 1-4	40.4	2,805	24.6	18.8	1,132
Primary 5-8	35.9	3,314	34.1	27.8	1,189
Secondary+	32.1	1,062	41.4	36.3	340
Wealth quintile					
Lowest	40.0	1,889	23.2	15.6	757
Second	41.2	2,170	26.3	22.2	894
Middle	37.6	2,206	25.5	19.4	830
Fourth	35.3	1,916	33.0	27.6	676
Highest	29.7	1,597	39.5	34.0	474
Total	37.1	9,777	28.4	22.7	3,630

Page 449, goal 5 – The numbers for condom use at last high-risk sex (population age 15-49) (%) are incorrect. The correct numbers are: Male 47.1, Female 30.1, and Total 37.9.