

ERRATA

National Statistical Service [Armenia], Ministry of Health [Armenia], and ORC Macro. 2006. *Armenia Demographic and Health Survey 2005*. Calverton, Maryland: National Statistical Service, Ministry of Health, and ORC Macro.

Pages 150 and 151 – Delete the two paragraphs on page 150 and Table 12.7 on page 151. Replace with the following text, figure, and table.

INFANT AND YOUNG CHILD FEEDING (IYCF) PRACTICES

Appropriate Infant and Young Child Feeding (IYCF) practices include breastfeeding through the age of two years, the introduction of solid and semisolid foods at age 6 months, and a gradual increase in the amount of food given and the frequency of feeding as the child gets older. The average, healthy **breastfed child** should receive solid and semisolid foods 2-3 times per day at age 6-8 months, and 3-4 times per day at age 9-23 months, with an additional snack 1-2 times per day. The minimum frequencies for feeding children in developing countries are based on the energy output of complementary foods. The energy needs of children are based on age-specific total daily energy requirements, plus 2 SD (to cover almost all children), minus the average energy intake from breast milk. Infants with low breast milk intake need to be fed more frequently than those with high breast milk intake. However, care should be taken that feeding frequencies do not exceed recommended input from complementary foods because excessive feeding can result in displacement of breast milk (PAHO/WHO, 2003).

Although the World Health Organization recommends that infants be breastfed up to the age of two years, some infants are not breastfed at all, or stopped breastfeeding before their second birthday. Guidelines have been developed for these children, who may not have been breastfed because their mother was HIV positive, or because their mother had died, or for other reasons (WHO, 2005). It is recommended that the **nonbreastfed child** be given solid and semisolid foods 4-5 times per day at age 6-23 months, with an additional snack 1-2 times per day.

Appropriate nutrition includes feeding children a variety of foods to ensure that nutrient requirements are met. Studies have shown that plant-based complementary foods by themselves are not sufficient to meet the needs of some children for certain micronutrients (WHO/UNICEF, 1998). Therefore, it is advised that children eat meat, poultry, fish, or eggs daily, or as often as possible. Vegetarian diets may not meet children's nutrient requirements unless supplements or fortified foods are also provided. Vitamin A-rich fruits and vegetables should be consumed daily, and the diets of children should include an adequate amount of fat. Fat is important in the diets of infants and young children because it provides essential fatty acids, facilitates absorption of fat-soluble vitamins (such as vitamin A), and enhances dietary energy density and palatability. Tea and coffee are not recommended for children because they contain compounds that inhibit iron absorption. Sugary drinks and excessive juice consumption should be avoided because other than energy they contribute little to the diet and decrease the child's appetite for more nutritious foods (PAHO/WHO, 2003).

In summary,


- Breastfed children age 6-23 months should receive animal-source foods and vitamin A-rich fruits and vegetables daily (PAHO/WHO, 2003). Because first foods almost always include a grain- or tuber-based staple, it is unlikely that young children who eat less than three food groups will receive both an animal-source food and a vitamin A-rich fruit or vegetable. Therefore, three food

groups are considered the minimum number appropriate for breastfed children (Arimond and Ruel, 2004).

- Breastfed infants age 6-8 months should receive complementary foods 2-3 times per day, with 1-2 snacks; breastfed children age 9-23 months should be receive meals 3-4 times per day, with 1-2 snacks (PAHO/WHO, 2003). Table 12.7 shows the percentage of breastfed children who were fed at least the minimum number of times per day for their age (i.e., twice for infants age 6-8 months and three times for children age 9-23 months).
- Nonbreastfed children age 6-23 months should receive milk or milk products to ensure that their calcium needs are met. In addition, they need animal-source foods and vitamin A-rich fruits and vegetables. Four food groups are considered the minimum number appropriate for nonbreastfed young children.
- Nonbreastfed children age 12-23 months should be fed meals 4-5 times per day, with 1-2 snacks (WHO, 2005). Table 12.7 shows the percentage of nonbreastfed children age 6-23 who were fed at least the minimum number of times per day (i.e., four times).

According to the results presented in Table 12.7, 96 percent of (youngest) Armenian children age 6-23 months living with the mother received breast milk or breast milk substitutes during the 24-hour period before the survey, 84 percent had an adequately diverse diet—i.e., they had been fed foods from the appropriate number of food groups, depending on their age and breastfeeding status—and 62 percent had been fed the minimum number of times appropriate for their age. Feeding practices for more than half of Armenian children age 6-23 months met the minimum standards with respect to all three of the IYCF feeding practices (see figure below).

Infant and Young Child Feeding (IYCF) Practices, Armenia 2005



Breastfed children are more likely than nonbreastfed children to be fed the minimum number of times per day but less likely to receive foods from the minimum number of food groups (for their age). Children age 9-11 months are more likely to meet the minimum standards than younger children. Children in urban areas (59 percent) are more likely to be fed according to the recommended guidelines, compared with the children in rural areas (49 percent). It is difficult to draw conclusions about regional differences in feeding practices of Armenian children because of the small sample size. Differences in feeding practices by mother's education and household wealth are also minimal.

Arimond, M., and M.T. Ruel. 2004. Dietary Diversity is Associated with Child Nutritional Status: Evidence from 11 Demographic and Health Surveys. *Journal of Nutrition* 134: 2579.

Pan-American Health Organization and World Health Organization (PAHO/WHO). 2003. *Guiding Principles for Complementary Feeding of the Breastfed Child*. Washington, D.C. and Geneva, Switzerland: PAHO/WHO.

World Health Organization (WHO). 2005. *Guiding Principles for Feeding Nonbreastfed Children 6 to 24 Months of Age*. Geneva, Switzerland: WHO.

World Health Organization and United Nations Children's Fund (WHO/UNICEF). 1998. *Complementary Feeding of Young Children in Developing Countries: A Review of Current Scientific Knowledge*. Geneva: World Health Organization, WHO/NUT98.1.

Table 12.7 Infant and young child feeding (IYCF) practices in Armenia

Percentage of youngest children age 6-23 months living with the mother who are fed according to three IYCF feeding practices based on the number of food groups received and the number of times child was fed during the past 24 hours (the day and night preceding the survey), by breastfeeding status and background characteristics, Armenia 2005

Background characteristic	Among breastfed children 6-23 months, percentage fed:				Among nonbreastfed children 6-23 months, percentage fed:				Among all children 6-23 months, percentage fed:					
	3+ food groups ¹	Minimum times or more ²	Both 3+ food groups and minimum times or more	Number of children (weighted)	Milk or milk products ³	4+ food groups	4+ times or more	With 3 IYCF practices ⁴	Number of children (weighted)	Breast milk or milk products	3+ or 4+ food groups ⁵	Minimum times or more ⁶	With all 3 IYCF practices	Number of children (weighted)
Age														
6-8	(47.9)	(66.6)	(43.7)	52	100.0	72.7	50.6	42.5	21	100.0	54.9	62.1	43.4	73
9-11	(90.0)	(76.1)	(76.1)	41	(97.3)	(81.4)	(66.4)	(51.2)	46	98.6	85.4	70.9	62.9	87
12-17	(85.5)	(62.6)	(59.5)	43	93.6	89.1	50.1	48.4	85	95.8	87.8	54.4	52.1	128
18-23	(95.0)	(68.3)	(68.3)	28	90.1	93.1	60.2	55.3	118	92.0	93.5	61.8	57.8	147
Sex														
Male	74.0	62.9	54.6	100	93.7	87.5	55.9	52.2	146	96.3	82.0	58.7	53.2	246
Female	79.9	76.4	68.6	65	92.6	89.2	59.1	50.6	124	95.1	86.0	65.1	56.7	188
Residence														
Urban	77.1	70.8	65.6	92	94.8	89.3	61.1	55.3	165	96.7	84.9	64.5	59.0	258
Rural	75.2	64.9	53.1	72	90.6	86.7	51.5	45.4	104	94.4	82.0	57.0	48.6	177
Region														
Yerevan	(81.0)	(76.6)	(71.9)	57	(98.6)	(93.3)	(65.5)	(60.4)	111	99.1	89.1	69.3	64.3	168
Aragatsotn	*	*	*	8	*	*	*	*	20	(95.6)	(94.6)	(54.5)	50.8	28
Ararat	*	*	*	22	(87.6)	(68.1)	(45.4)	33.6	21	(94.0)	(60.3)	(50.1)	33.4	43
Armavir	*	*	*	11	*	*	*	*	27	(98.8)	(93.7)	(56.1)	51.6	39
Gegharkunik	*	*	*	18	*	*	*	*	16	(95.1)	(79.7)	(59.7)	51.7	33
Lori	*	*	*	9	*	*	*	*	15	*	*	*	*	24
Kotayk	*	*	*	15	*	*	*	*	22	(95.3)	(83.4)	(40.9)	35.1	36
Shirak	*	*	*	6	*	*	*	*	21	(75.1)	(49.4)	(53.1)	41.6	27
Syunik	*	*	*	7	*	*	*	*	9	(100.0)	(84.2)	(71.8)	64.8	17
Vayots Dzor	*	*	*	2	*	*	*	*	3	*	*	*	*	5
Tavush	*	*	*	10	*	*	*	*	5	(91.2)	(96.1)	(70.7)	65.9	15
Mother's education														
Primary	*	*	*	0	*	*	*	*	0	*	*	*	*	6
Secondary	80.1	67.4	58.8	124	91.5	85.5	55.9	48.8	206	94.7	83.5	60.2	52.5	330
More than secondary	(64.8)	(70.7)	(64.0)	(41)	(98.6)	(97.0)	(68.3)	(66.1)	58	99.2	83.7	69.3	65.2	99
Wealth quintile														
Lowest	(75.4)	(63.0)	(48.7)	28	(92.9)	(92.6)	(59.0)	(51.8)	52	95.4	86.5	60.4	50.8	80
Second	(87.4)	(81.8)	(76.0)	38	87.7	77.0	44.2	41.2	59	92.5	81.1	58.9	54.8	97
Middle	(60.3)	(56.0)	(45.3)	37	(87.9)	(76.0)	(57.3)	(43.2)	40	93.7	68.5	56.7	44.3	76
Fourth	(78.6)	(63.6)	(55.7)	42	96.7	97.6	68.1	65.2	66	98.0	90.2	66.4	61.5	107
Highest	*	*	*	21	(99.1)	(94.4)	(57.4)	(51.8)	53	99.4	90.6	63.9	59.9	74
Total	76.3	68.2	60.1	165	93.2	88.3	57.4	51.5	270	95.8	83.7	61.5	54.7	435

Note: Figures in parentheses are based on 25-49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

¹ Food groups: a) infant formula, milk other than breast milk, cheese or yogurt or other milk products; b) foods made from grains, roots, and tubers, including porridge and, fortified baby food from grains; c) vitamin A-rich fruits and vegetables (and red palm oil); d) other fruits and vegetables; e) eggs; f) meat, poultry, fish, and shellfish (and organ meats); g) legumes and nuts; h) foods made with oil, fat, butter.

² At least twice a day for breastfed infants age 6-8 months and at least three times a day for breastfed children age 9-23 months

³ Includes commercial infant formula, fresh, tinned and powdered animal milk, and cheese, yogurt and other milk products

⁴ Nonbreastfed children age 6-23 months are considered to be fed with three IYCF practices if they receive other milk or milk products and are fed at least the minimum number of food groups and the minimum number of times per day.

⁵ 3+ food groups for breastfed children and 4+ food groups for nonbreastfed children

⁶ Fed solid or semisolid food at least twice a day for infants age 6-8 months, 3+ times a day for other breastfed children, and 4+ times a day for nonbreastfed children