

ERRATA

in

Riese, Sara, Shireen Assaf, and Thomas Pullum. 2021. *Measurement Approaches for Effective Coverage Estimation*. DHS Methodological Reports No. 31. Rockville, Maryland, USA: ICF.

| Date of correction | Correction |
|---------------------------|--|
| Nov 15, 2021 | Page 15, Table 6 New data are highlighted in tables below. |
| | Page 17, Figure 2 Confidence intervals of quality-adjusted coverage in Approaches 2 and 3 corrected. |
| | Page 18, Figure 3 Confidence intervals of intervention-adjusted and quality-adjusted coverage in Approaches 2 and 3 corrected. |
| | Page 19, Figure 4 Confidence intervals of quality-adjusted coverage in Approaches 2 and 3 corrected. |
| | Page 20, Figure 5 Confidence intervals of intervention-adjusted and quality-adjusted coverage in Approaches 2 and 3 corrected. |
| | Appendix A, Table 4 Confidence intervals of client-level process quality measures were added and are highlighted in tables below. |

The calculation of confidence intervals for client-level measures from the two facility census SPAs (Haiti and Malawi) was corrected. This correction resulted in changes to the confidence intervals in Table 6 on page 15, Figures 2-5, as well as in Appendix A Table 4. No text was changed. These changes do not affect the interpretation of the estimates or the overall findings of the report.

Table 6 Values of each component of each effective coverage measurement approach

| | Coverage | Readiness | Receipt of complete intervention | Process quality |
|-------------------|-------------------|-------------------|----------------------------------|-------------------|
| Haiti | | | | |
| ANC | | | | |
| Approach 1 | 0.91 [0.90, 0.93] | NA | 0.67 [0.64, 0.69] | 0.84 [0.83, 0.85] |
| Approach 2 | 0.91 [0.90, 0.93] | 0.70 [0.70, 0.70] | 0.67 [0.64, 0.69] | 0.59 [0.57, 0.60] |
| Approach 3 | 0.91 [0.90, 0.93] | 0.58 [0.58, 0.58] | 0.67 [0.64, 0.69] | 0.55 [0.53, 0.56] |
| Sick child | | | | |
| Approach 1 | 0.26 [0.23, 0.29] | NA | 0.20 [0.17, 0.23] | NA |
| Approach 2 | 0.26 [0.23, 0.29] | 0.73 [0.73, 0.73] | 0.45 [0.40, 0.50] | 0.46 [0.43, 0.49] |
| Approach 3 | 0.26 [0.23, 0.29] | 0.59 [0.59, 0.59] | 0.45 [0.40, 0.50] | 0.37 [0.36, 0.39] |
| Malawi | | | | |
| ANC | | | | |
| Approach 1 | 0.98 [0.98, 0.99] | NA | 0.51 [0.49, 0.52] | 0.70 [0.69, 0.71] |
| Approach 2 | 0.98 [0.98, 0.99] | 0.50 [0.50, 0.50] | 0.51 [0.49, 0.52] | 0.49 [0.47, 0.51] |
| Approach 3 | 0.98 [0.98, 0.99] | 0.62 [0.62, 0.62] | 0.51 [0.49, 0.52] | 0.57 [0.55, 0.58] |
| Sick child | | | | |
| Approach 1 | 0.53 [0.50, 0.55] | NA | 0.42 [0.40, 0.44] | NA |
| Approach 2 | 0.53 [0.50, 0.55] | 0.85 [0.85, 0.85] | 0.92 [0.89, 0.94] | 0.29 [0.27, 0.32] |
| Approach 3 | 0.53 [0.50, 0.55] | 0.64 [0.64, 0.64] | 0.92 [0.89, 0.94] | 0.33 [0.31, 0.35] |

Figure 2 ANC effective coverage cascade in Haiti for each measurement approach




Figure 3 Sick child effective coverage cascade (among children with ARI/diarrhea symptoms in last 2 weeks) in Haiti for each measurement approach




Figure 4 ANC effective coverage cascade in Malawi for each measurement approach





Figure 5 Sick child effective coverage cascade (among children with ARI/diarrhea symptoms in last 2 weeks) in Malawi for each measurement approach



Appendix Table 4 ANC process indicators among women attending their first visit and sick child intervention and process indicators

| | Haiti | Malawi | Nepal | Senegal | Tanzania |
|--|------------------|-------------------|------------------|------------------|------------------|
| ANC process quality | | | | | |
| Client history | 98.7 [97.6,99.3] | 96.1 [94.4,97.3] | 87.3 [81.2,91.6] | 96.2 [92.2,98.2] | 97.2 [95.9,98.2] |
| Urine test | 66.0 [60.4,71.1] | 4.3 [2.8,6.6] | 38.9 [31.9,46.5] | 68.0 [51.3,81.1] | 41.4 [36.3,46.8] |
| Hb test | 70.9 [65.4,75.9] | 9.6 [7.4,12.5] | 41.2 [34.1,48.6] | 40.7 [29.8,52.7] | 46.6 [41.2,52.1] |
| Syphilis test | 64.0 [58.1,69.5] | 12.6 [10.1,15.8] | 22.8 [17.8,28.7] | 64.6 [48.7,77.8] | 41.9 [36.8,47.2] |
| Checked blood pressure | 95.8 [92.9,97.5] | 48.0 [44.2,51.8] | 83.4 [76.7,88.5] | 97.6 [93.9,99.1] | 72.2 [66.6,77.2] |
| Measured weight | 91.4 [87.6,94.1] | 73.7 [70.0,77.0] | 72.3 [63.9,79.5] | 97.8 [93.8,99.2] | 79.2 [73.4,83.9] |
| Checked fundal height | 70.5 [65.1,75.3] | 94.9 [93.2,96.2] | 39.1 [32.9,45.7] | 84.8 [61.1,95.2] | 94.8 [92.7,96.2] |
| Physical examination/vaginal exam | 10.4 [7.3,14.6] | 3.5 [2.5,4.9] | 2.8 [1.4,5.5] | 74.8 [56.8,87.0] | 10.3 [7.8,13.6] |
| Checked edema | 18.6 [14.7,23.3] | 64.8 [61.0,68.3] | 26.2 [20.8,32.4] | 78.5 [68.0,86.2] | 58.6 [54.1,63.0] |
| Checked client card | 43.9 [37.6,50.4] | 99.9 [99.2,100.0] | 90.2 [84.2,94.1] | 86.4 [78.3,91.8] | 94.1 [91.9,95.8] |
| Examined palms for anemia/check pallor for anemia | 91.4 [87.6,94.1] | 73.7 [70.0,77.0] | 72.3 [63.9,79.5] | 97.8 [93.8,99.2] | 79.2 [73.4,83.9] |
| Daily oral iron and folic acid supplementation (counseled or prescribed) | 80.6 [76.2,84.4] | 90.7 [88.3,92.6] | 77.7 [71.7,82.8] | 97.6 [95.3,98.8] | 86.5 [83.0,89.5] |
| Tetanus toxoid vaccination | 29.7 [24.3,35.8] | 45.2 [41.4,49.0] | 31.3 [25.4,37.9] | 66.3 [54.6,76.3] | 52.1 [47.4,56.8] |
| Danger signs of current pregnancy | 75.3 [70.5,79.5] | 62.5 [58.6,66.3] | 58.0 [50.2,65.5] | 74.0 [55.9,86.4] | 74.3 [69.0,78.9] |
| Counseling on healthy eating | 49.2 [43.5,55.0] | 44.5 [40.6,48.3] | 52.9 [45.6,60.2] | 61.2 [47.9,73.0] | 34.7 [30.2,39.5] |
| Informed the client about the progress of the pregnancy | 21.5 [17.6,26.0] | 56.7 [52.9,60.4] | 24.3 [18.5,31.1] | 27.9 [19.6,38.0] | 49.9 [45.2,54.7] |
| Counseled on breastfeeding | 4.3 [2.4,7.5] | 9.9 [7.5,12.9] | 2.1 [1.0,4.6] | 4.8 [2.4,9.5] | 15.8 [13.0,19.1] |
| Wrote on client card | 38.9 [32.7,45.4] | 99.9 [99.2,100.0] | 88.5 [82.5,92.7] | 86.1 [78.0,91.6] | 94.1 [91.9,95.7] |
| Encouraged questions | 66.2 [60.1,71.9] | 78.0 [74.6,81.1] | 37.1 [30.2,44.4] | 65.2 [49.6,78.2] | 79.5 [74.9,83.4] |
| Sick child complete intervention coverage | | | | | |
| Child diagnosed with pneumonia | 2.8 [2.1,3.8] | 11.6 [10.3,13.1] | 8.2 [6.5,10.2] | 5.0 [3.0,8.1] | 11.6 [10.2,13.1] |
| Child diagnosed with diarrhea | 15.5 [13.7,17.5] | 10.4 [9.2,11.7] | 15.8 [13.8,18.0] | 29.7 [14.0,52.2] | 17.2 [15.5,19.0] |
| Child with pneumonia given any antibiotic | 62.4 [48.4,74.5] | 98.0 [95.9,99.1] | 86.7 [79.1,91.8] | 94.6 [80.5,98.7] | 96.4 [94.0,97.8] |
| Child with diarrhea given ORS or zinc | 41.7 [36.3,47.2] | 85.9 [81.7,89.2] | 74.4 [67.7,80.1] | 82.1 [59.9,93.4] | 64.9 [60.0,69.5] |
| Child with any of these 2 given treatment | 44.9 [39.8,50.1] | 91.6 [89.1,93.5] | 77.8 [72.4,82.4] | 84.4 [66.9,93.6] | 76.9 [73.5,80.0] |

Continued...

Appendix Table 4—continued

| | Haiti | Malawi | Nepal | Senegal | Tanzania |
|---|------------------|------------------|------------------|------------------|------------------|
| Sick child process quality | | | | | |
| Provider counted respiration for 60 seconds | 37.7 [32.9,42.7] | 24.2 [22.0,26.5] | 54.8 [49.1,60.3] | 69.4 [54.3,81.2] | 22.5 [19.4,26.0] |
| Provider weighed client | 81.3 [77.7,84.5] | 26.5 [24.3,28.7] | 50.2 [45.5,54.9] | 61.7 [38.6,80.4] | 16.1 [13.7,18.8] |
| Provider plotted weight on growth chart | 10.5 [8.4,13.1] | 7.3 [6.0,8.7] | 16.7 [13.6,20.3] | 96.6 [94.1,98.0] | 4.7 [3.3,6.5] |
| Provider checked palms/conjunctiva for pallor | 49.4 [45.9,52.8] | 43.6 [41.5,45.9] | 14.5 [12.3,17.0] | 17.3 [10.4,27.5] | 41.7 [38.7,44.8] |
| Provider discussed weight/growth/growth chart | 9.7 [7.9,11.8] | 8.7 [7.6,9.8] | 4.1 [3.0,5.6] | 11.5 [6.8,18.6] | 24.3 [21.8,27.0] |
| Checked skin turgor for dehydration | 55.0 [49.0,60.8] | 42.1 [37.8,46.4] | 29.1 [23.5,35.5] | 69.6 [43.6,87.1] | 46.0 [41.5,50.6] |
| Provider took temperature | 85.3 [82.3,87.8] | 42.1 [39.9,44.2] | 47.6 [43.6,51.7] | 96.1 [92.9,97.9] | 42.5 [38.8,46.2] |
| Provider checked for edema | 5.8 [4.6,7.3] | 8.0 [6.9,9.2] | 1.6 [1.0,2.6] | 8.0 [5.0,12.4] | 6.0 [4.7,7.6] |
| Felt the child for fever or body hotness | 16.1 [13.8,18.6] | 42.3 [40.0,44.5] | 29.0 [25.8,32.3] | 5.5 [3.2,9.3] | 33.9 [31.0,37.0] |
| Checked for neck stiffness | 3.3 [2.4,4.4] | 1.9 [1.5,2.6] | 0.9 [0.5,1.5] | 0.6 [0.2,1.5] | 5.8 [4.6,7.4] |
| Looked in child's ear | 16.1 [13.5,19.1] | 4.0 [3.3,4.9] | 10.1 [8.1,12.6] | 10.8 [6.2,18.2] | 12.7 [10.9,14.8] |
| Felt behind child's ear | 11.3 [9.6,13.3] | 4.4 [3.6,5.5] | 5.8 [4.5,7.4] | 9.4 [4.9,17.5] | 9.0 [7.4,10.9] |
| Checked vaccination status | 21.3 [18.9,23.9] | 16.5 [15.1,18.0] | 5.9 [4.8,7.4] | 60.0 [44.4,73.9] | 24.3 [21.8,26.9] |
| Check mouth and throat | 19.2 [16.7,21.9] | 9.6 [8.4,11.0] | 11.4 [9.6,13.5] | 23.0 [8.5,49.2] | 17.1 [15.0,19.3] |
| Provider asked / caretaker mentioned if child unable to drink or breastfeed | 19.5 [17.3,21.9] | 27.8 [25.9,29.9] | 21.6 [18.9,24.5] | 2.4 [1.2,4.6] | 27.8 [25.3,30.3] |
| Provider asked / caretaker mentioned cough or difficult breathing | 64.5 [61.9,67.1] | 73.8 [71.8,75.6] | 55.9 [52.4,59.3] | 59.8 [40.5,76.5] | 75.4 [73.1,77.5] |
| Provider asked / caretaker mentioned diarrhea | 32.0 [29.1,35.1] | 39.0 [36.9,41.1] | 41.2 [38.1,44.5] | 42.8 [25.6,62.0] | 59.4 [56.6,62.1] |
| Provider asked / caretaker mentioned fever | 65.1 [62.2,68.0] | 77.3 [75.2,79.2] | 77.1 [73.9,80.0] | 66.5 [45.4,82.6] | 93.3 [92.1,94.3] |
| Provider asked / caretaker mentioned convulsions | 3.7 [2.9,4.8] | 9.3 [8.1,10.6] | 4.5 [3.3,6.3] | 0.8 [0.4,2.0] | 20.1 [17.7,22.7] |
| Provider asked caretaker about vomiting | 24.7 [22.4,27.2] | 34.8 [32.8,36.9] | 20.4 [17.7,23.5] | 16.7 [11.1,24.3] | 48.7 [45.7,51.7] |
| Provider asked / caretaker mentioned ear pain | 4.9 [3.9,6.0] | 4.8 [3.9,5.8] | 17.6 [14.8,20.8] | 3.6 [1.6,8.0] | 16.4 [14.1,19.0] |
| Provider asked about normal feeding/breastfeeding when not ill | 39.9 [36.7,43.1] | 14.2 [12.8,15.8] | 17.7 [15.1,20.7] | 26.7 [17.9,37.8] | 27.9 [25.4,30.5] |
| Provider asked about feeding or breastfeeding during this illness | 31.5 [28.9,34.1] | 15.5 [14.0,17.1] | 16.4 [14.0,19.2] | 7.8 [4.7,12.7] | 18.8 [16.8,21.0] |
| Provider general information about feeding/breastfeeding | 27.9 [25.0,30.9] | 8.7 [7.5,9.9] | 11.1 [9.2,13.3] | 25.4 [17.1,35.9] | 14.4 [12.6,16.4] |
| Advise extra fluids during this sickness | 16.8 [14.6,19.2] | 7.5 [6.5,8.6] | 17.5 [15.0,20.3] | 6.8 [4.1,10.9] | 11.9 [10.4,13.7] |
| Advise continued feeding during sickness | 15.2 [13.2,17.5] | 11.8 [10.5,13.1] | 16.7 [14.4,19.3] | 9.5 [6.1,14.7] | 13.1 [11.4,14.9] |
| Provider described more than 1 danger sign requiring return to facility | 5.8 [4.6,7.3] | 11.3 [10.1,12.7] | 7.1 [5.7,8.7] | 6.3 [3.8,10.1] | 16.6 [14.4,19.1] |
| Provider stated diagnosis to caretaker | 15.2 [13.2,17.4] | 40.5 [38.4,42.7] | 34.9 [31.1,39.0] | 12.1 [7.8,18.2] | 48.8 [45.4,52.2] |
| Provider explained dosing if medication prescribed | 73.0 [69.2,76.4] | 51.1 [48.7,53.5] | 56.2 [51.6,60.6] | 66.8 [51.8,79.0] | 50.5 [46.4,54.7] |
| Provider asked caretaker to repeat the instructions for giving medicines at home (if medication prescribed) | 5.9 [4.6,7.6] | 5.9 [5.0,7.0] | 3.7 [2.4,5.5] | 2.0 [1.1,3.8] | 11.2 [9.0,13.9] |
| Provider discussed follow-up visit | 35.3 [31.8,38.9] | 20.7 [19.1,22.3] | 25.9 [22.8,29.2] | 60.3 [45.1,73.8] | 22.8 [20.3,25.5] |
| Provider used visual aids | 0.3 [0.1,0.9] | 0.2 [0.1,0.4] | 0.2 [0.1,0.7] | 1.5 [0.6,3.2] | 0.9 [0.4,1.8] |
| Provider recorded on child's health card/booklet | 17.8 [14.8,21.1] | 87.8 [86.1,89.4] | 65.3 [60.3,70.1] | 21.4 [7.2,48.9] | 21.5 [18.5,24.8] |